

VPARP antibody to the minor vault protein, clone P193-6 Monoclonal Antibody

Catalog No.: MON 9040

Quantity: 1 ml

Specificity

p193-6 reacts with an internal epitope (amino acids 593-599, FSKVEDY) of the minor vault protein (p193 or VPARP), which is overexpressed in various human non-P-glycoprotein MDR tumor cell lines, accordingly to an increase in the number of vault particles. p193-6 was raised against an *E. coli* lysate transformed with the pET28a(+) expression vector containing amino acids 408-611 of the p193 cDNA.

Immunoglobulin type

Murine IgG_{2b} kappa

Use

p193 expression reflects vault density in human normal tissue and tumor samples. The antibody can be diluted 1:10-20 for immunohistochemical staining of paraformaldehyde (4%) fixed cytospin preparations and frozen tissue sections, and Western blotting (preferably with ECL). For immunohistochemistry apply the following pretreatment: 10 min 20 mM glycine in PBS (pH 7.5) and 10 min 6 N guanidine hydrochloride in 50 mM Tris-HCl (pH 7.5); Schroeijers *et al.*, 2000. Optimal conditions for staining paraffin-embedded tissues and FACS analyses have not yet been defined

Presentation

1 ml vials (>200 tests) containing antibody dissolved in 1% BSA containing 0.1% sodium azide. Concentration approx. 250 µg immunoglobulin/ml.


Storage

4°C. For extended storage, freeze at -20°C.

Literature


Schroeijers AB et al., Cancer Res. 60: 1104, 2000; Kickhoefer VA et al., J. Cell Biol. 146: 917, 1999; Kickhoefer VA et al., J. Bio. Chem. 273: 8971, 1998; Scheper RJ et al., Cancer Res. 53: 1475, 1993; Scheffer GL et al, Nature Med. 1: 578, 1995; Izquierdo MA et al., Cytotechnology 27: 137, 1998.

FOR RESEARCH USE ONLY, NOT FOR DRUG, DIAGNOSTIC OR OTHER USE.



Toll Free: 888-769-1246 E-mail: info@cellsciences.com
Web Site: www.cellsciences.com
Fax: 781-828-0542

cellsciences.com



Toll Free: 888-769-1246 E-mail: info@cellsciences.com
Web Site: www.cellsciences.com
Fax: 781-828-0542