

## TNFRSF1B

### Mouse Anti-Human TNFR2 Clone 80M2 mAb

<b>Catalog No.</b>	MON9059	<b>Quantity:</b>	100 µg
<b>Alternate Names:</b>	p75, TBPII, TNFBR, TNFR2, CD120b, TNFR1B, TNFR80, TNF-R75, p75Tnfr, TNF-R-II		
<b>Gene ID:</b>	7133		
<b>Specificity:</b>	The antibody reacts with the extra-cellular part of the TNF-RII. It also reacts with the soluble receptor. TNF-RII is present on most cell types and is considered to be an important co-stimulator together with TNF-RI in cellular responses to TNF-alpha, in particular the transmembrane form of TNF-alpha. A dominant role of TNF-RII has been shown in thymocyte activation by TNF-alpha		
<b>Clone:</b>	80M2		
<b>Immunoglobulin Type:</b>	Mouse IgG		
<b>Clone:</b>	80M2		
<b>Use:</b>	The antibody is a <u>non-agonistic receptor modulating</u> antibody. It enhances in vitro TNF responses by increasing the affinity of the soluble form to TNF-alpha to TNF-RII. The antibody is useful for flow cytometry and immunohistology on frozen sections. Furthermore the antibody is useful for immuno precipitation, Western blotting and immuno assays. The reactivity of the antibody with cellbound TNF-Receptor is minimally inhibited by high concentrations of TNF-alpha.		
<b>Instructions for Use:</b>	For immunohistology, flow cytometry and Western blotting dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:10. Advised positive controls for frozen sections are human lymphnodes and for flow cytometry PHA activated T cells.		
<b>Presentation:</b>	1 ml (100 µg/ml) 0.2 µm filtered antibody solution in PBS, containing 0.02% sodium azide and 0.1% bovine serum albumin		
<b>Storage &amp; Stability:</b>	Product should be stored at 2-4°C. Under recommended storage conditions, product is stable for one year.		

**NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.**