

## IL10, Affinity Purified Human pAb

<b>Catalog No.</b>	PA0602	<b>Quantity:</b>	100 µg
<b>Description:</b>	Rabbit Anti-human IL10 Affinity Purified polyclonal antibody		
<b>Specificity:</b>	Human		
<b>Host:</b>	Rabbit		
<b>Immunogen:</b>	Recombinant Human IL-10		
<b>Formulation:</b>	A sterile filtered antibody solution (1.0 mg/ml) was lyophilized from 0.5X PBS pH 7.4.		
<b>Purification:</b>	Affinity Chromatography		
<b>Reconstitution:</b>	<b>Centrifuge vial prior to opening.</b> Reconstitute in sterile water to a concentration of ≤0.2 mg/ml.		
<b>Storage &amp; Stability:</b>	Stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted according to the above directions, the antibody is stable for at least six weeks at 2-4°C. <b>Avoid repeated freeze-thaw cycles.</b>		
<b>Applications:</b>	<p><b>Neutralization:</b> To yield one-half maximal inhibition [ND50] of the biological activity of human IL-10 (30 ng/ml), a concentration of 4.0-6.0 µg/ml of this antibody is required.</p> <p><b>ELISA:</b> To detect human IL-10 by direct ELISA (using 100 µl/well antibody solution) a concentration of at least 0.5 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2-0.4 ng/well of recombinant human IL-10.</p> <p><b>Western Blot:</b> To detect human IL-10 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant human IL-10 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.</p> <p>The optimal concentration should be determined by the user for each specific application.</p>		

**NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.**

