

## Anti-human LYVE-1 Polyclonal Antibody, Protein A Purified

Catalog Number: PA0843A Size: 100 μg

PA0843B **Size**: 200 μg

**Description**: Produced from sera of rabbits immunized with highly pure recombinant human

soluble LYVE-1 produced in insect cells. The recombinant soluble LYVE-1 consists of amino acid 24 (Ser) to 232 (Gly) and is fused to a C-terminal His-tag (6xHis). LYVE-1 has been identified as a major receptor for HA (extracellular matrix glycosaminoglycan hyaluronan) on the lymph vessel wall. The deduced amino acid sequence of LYVE-1 predicts a 322-residue type I integral membrane polypeptide 41% similar to the CD44 HA receptor with a 212-residue extracellular domain containing a single Link module the prototypic HA binding domain of the Link protein superfamily. Like CD44, the LYVE-1 molecule binds both soluble and immobilized HA. However, unlike CD44, the LYVE-1 molecule co localizes with HA on the luminal face of the lymph vessel wall and is completely absent from blood vessels. Hence, LYVE-1 is the first lymph-specific HA receptor to be characterized and is a uniquely powerful marker for lymph vessels themselves.

Host Species: Rabbits

Antigen: Recombinant human soluble Lyve-1

**Purification:** Protein-A Chromatography (+his tag depleted)

Stabilizer: none

**Buffer:** lyophilized from PBS, pH 7.4 w/o preservative

Formulation: lyophilized rabbit IgG

**Reconstitution**: The lyophilized IgG is stable at 4°C for at least one month and for greater than a

year when kept at  $-20^{\circ}$ C. Centrifuge vial before opening to bring contents to the bottom. When reconstituted in sterile water/PBS to a concentration of greater than 0.5 mg/ml the antibody is stable for at least six weeks at 2-4°C. **Avoid** 

Toll Free: 888-769-1246

Phone: 781-828-0610

Fax: 781-828-0542

E-mail: info@cellsciences.com

Web Site: www.cellsciences.com

repeated freeze-thaw cycles.

**Applications** 

**ELISA:** Use at 1-15  $\mu$ g/ml.

**Western Analysis:** Use at a concentration of 1-2 µg/ml with the appropriate secondary reagents.

Immunohistochemistry: Use at 6-30 µg/ml.

Optimal dilutions should be determined by each laboratory for each application.

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.