

Goat Anti-Human Immunoglobulin G + Immunoglobulin A + Immunoglobulin M Peroxidase pAb

| | | | |
|--------------------------|--|------------------|--------|
| Catalog No. | PS103P | Quantity: | 1.2 mg |
| Description: | Antibodies were raised in goat to human IgG, IgM and IgA. Specific antibodies were absorbed by incubation with Sepharose-bound human immunoglobulins. Specific antibodies were eluted by acidic buffer at pH 2.5 followed by neutralization and dialysis. After repeated binding with immobilized human Immunoglobulines a minimum of 65% protein bound. The purified polyclonal was conjugated to horseradish peroxidase according to the periodate method followed by gel-filtration and ion-exchange chromatography to clear unbound conjugate. | | |
| Specificity: | Human | | |
| Host: | Goat | | |
| Immunogen: | IgG, IgM and IgG. | | |
| Formulation: | 1 ml purified goat antiserum at 1.2 mg/ml protein in 50% glycerol with 0.1 M NaCl, 0.01M sodium phosphate at pH 7.5. | | |
| Purification: | Affinity Purified | | |
| Cross-Reactivity: | No cross reaction with albumin fraction of human serum other than IgG, IgM and IgA. No cross reaction with albumin fraction of other animal species. | | |
| Applications: | Recommended dilution as secondary antibody in ELISA, 1:1000. Recommended dilution as secondary antibody in dot-blotting, 1:100. | | |

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
480 Neponset Street
Bldg 12A
Canton, MA 02021

Toll Free: 888-769-1246
Phone: 781-828-0610
Fax: 781-828-0542

E-mail: info@cellsciences.com
Website: www.cellsciences.com