

IL17A

Mouse Anti-Human IL-17A (Clone B-C49) mAb, Azide Free

Catalog No.	CDM120A CDM120B CDM120C	Quantity:	200 µg 500 µg 1.0 mg
Alternate Names:	Interleukin-17A, Cytotoxic T-lymphocyte-associated antigen 8, CTLA-8		
Description:	Interleukin-17 (IL-17, or IL-17A) is the original member of the IL-17 family of cytokines. IL-17A is involved in inducing and mediating proinflammatory responses, commonly associated with allergic responses and induces the production of many other cytokines (such as IL-6, G-CSF, GM-CSF, IL-1β, TGF-β, TNF-α), chemokines (including IL-8, GRO-α and MCP-1) and prostaglandins (e.g. PGE2) from many cell types (fibroblasts, endothelial cells, epithelial cells, keratinocytes and macrophages). IL-17A function is also essential to a subset of CD4+ T-Cells called T helper 17 (Th17) cells. High levels of this cytokine are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis.		
UniProt ID:	Q16552		
Gene ID:	3605		
Concentration:	1.0 mg/ml		
Specificity:	Recognizes both Recombinant and Native Human IL-17A		
Hybridoma:	Myeloma X63/AG.8653 x BALB/c lymph node cells		
Immunogen:	Recombinant Human IL-17A		
Isotype:	Mouse IgG2bk		
Clone:	B-C49		
Formulation:	Sterile filtered PBS, carrier and preservative free.		
Purification:	Ion exchange chromatography		
Application Notes:	<p>ELISA: Use as a Capture Antibody at 1-5 µg/ml in a Human IL-17A sandwich ELISA in combination with Biotinylated Human IL-17A Clone B-B51 Detection Antibody (Cat No CDM432).</p> <p>ELISPOT: Use as a Capture Antibody at 5-10 µg/ml.</p> <p>The suggested coating concentration range should be optimized by each laboratory for each application.</p>		
Storage & Stability:	<p>Store at 2-8°C for up to 1 year or in working aliquots at -20°C to -80°C.</p> <p>Avoid repeated freeze-thaw cycles.</p>		

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.