

## Mouse IL-5 ELISA antibody pair

Catalog No. **CKM094**

**10-plate format**

### Coating antibodies (2 vials)

**Product:** Monoclonal antibody to mouse interleukin-5 (IL-5)  
**Isotype:** Rat IgG<sub>1</sub>  
**Production:** *In vitro* using serum free medium  
**Purification:** Ion exchange chromatography  
**Contents:** Each vial contains sufficient material for coating of five 96-well ELISPOT plates.  
**Buffer:** Prior to lyophilization: 0.25 ml PBS + 125 mM trehalose  
**Application:** Coating antibody in an ELISA system.  
**Reconstitution:** Inject 0.25 ml distilled water into the vial and dilute 100 times in PBS. The content of one vial is sufficient for five 96-well ELISA plates (480 determinations; 50 µl/well).

### Detection antibodies (2 vials)

**Product:** Biotinylated monoclonal antibody to mouse interleukin 5 (IL-5)  
**Isotype:** Rat IgG<sub>2a</sub>  
**Production:** *In vitro* using serum free medium  
**Purification:** Protein G-affinity chromatography  
**Labeling:** With Biotin-7-NHS (N-hydroxysuccinimide)  
**Buffer:** Prior to lyophilization: 0.5 ml PBS + 1% BSA + 125 mM trehalose  
**Application:** Inject 0.5 ml distilled water into the vial and dilute 100 times in PBS + 0.5% BSA + 0.05% Tween-20. The content of one vial is sufficient for five 96-well ELISA plates (480 determinations; 100 µl/well).

### Standards (5 vials)

**Product:** Recombinant mouse interleukin 5 (IL-5)  
**Application:** Cytokine standard in an ELISA system  
**Reconstitution:** Inject 0.5 ml distilled water into the vial. Use immediately.

### Conjugate (2 vials)

**Product:** SPP conjugate (Streptavidin-HRP)  
**Application:** Inject 0.5 ml distilled water into the vial and dilute 100 times in PBS + 0.5% BSA + 0.05% Tween-20. The content of one vial is sufficient for five 96-well ELISA plates (480 determinations; 100 µl/well).

### General

**Sensitivity:** < 5 pg/ml  
**Specificity:** Validated for detecting natural and recombinant mouse IL-5  
**Sterility:** Membrane filtered (0.2 µm)  
**Stability:** Lyophilized SPP conjugate is stable for at least one year at -20°C, the other lyophilized products are stable for at least one year at 4°C.  
After reconstitution, the antibodies are stable for one year at 4°C (if kept sterile) and SPP for minimal one year at -20°C. The reconstituted standard preparation should be used immediately.

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



**Cell Sciences®**  
Neponset Valley Tech Park  
480 Neponset St., Bldg. 12A  
Canton, MA 02021

Toll Free: 888 769-1246  
Phone: 781 828-0610  
Fax: 781 828-0542

E-mail: [info@cellsciences.com](mailto:info@cellsciences.com)  
Web Site: [www.cellsciences.com](http://www.cellsciences.com)