


BDNF

Recombinant Human Brain-Derived Neurotrophic Factor

Catalog No.	CRB600A CRB600B CRB600C CRB600D	Quantity:	2 µg 10 µg 1.0 mg 100 µg
Alternate Names:	BDNF, Abrineurin, ANON2, BULN2		
Description:	Brain Derived Neurotrophic Factor, or BDNF, is a nerve growth factor that supports neuron growth and survival. BDNF shares identical domains with two other neurotrophic factors known as, β -NGF and NT-3 (neurotrophin-3). BDNF binds with low affinity to a receptor known as LNGFR, which also binds NGF and NT-3, but mediates survival function by signaling through a high affinity receptor known as gp145/TrkB. Human, mouse, rat and pig BDNF are all cross-reactive.		
Gene ID:	627		
UniProt ID:	P23560		
Source:	<i>E. coli</i>		
Molecular Weight:	Homodimer (non-covalent), 13.6/27.3 kDa (120/240 aa)		
Formulation:	Lyophilized from a sterile filtered solution containing 0.1% Trifluoroacetic Acid (TFA).		
Purity:	> 95% by reducing and non-reducing SDS-PAGE		
Endotoxin Level:	≤ 1 EU/µg by kinetic LAL		
Biological Activity:	ED ₅₀ ≤ 2 µg/ml, as determined by the dose dependent proliferation of C6 cells.		
Specific Activity:	$\geq 5.0 \times 10^2$ units/mg		
Amino Acid Sequence:	MHSDPARRGE LSVCDISEW VTAADKKTAV DMSGGTVTVL EKVPVSKGQL KQYFYETKCN PMGYTKEGCR GIDKRHWNSQ CRTTQSYVRA LTMDSKKRIG WRFIRIDTSC VCTLTIKRGR		
Reconstitution:	Centrifuge vial prior to opening. Add sterile distilled water to a recommended concentration of 0.1 mg/mL and gently pipet solution up and down sides of vial. DO NOT VORTEX. Allow several minutes for complete reconstitution.		
Storage & Stability:	Store as supplied at -20°C to -80°C for up to 1 year. Upon reconstitution, prepare working aliquots and store at -20°C to -80°C. It is recommended that a carrier protein such as 0.1% HSA or BSA is added for long term storage. Avoid repeated freeze-thaw cycles.		





Human BDNF Gel

Figure: 1 ug in each lane (-) non-reducing conditions and (+) reducing conditions in a 4-20% Tris-Glycine gel, stained with Coomassie Blue. Human BDNF is a non-covalently linked homodimer and has a predicted MW of 27.3 kDa.

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.