


NTN1

Recombinant Human Netrin-1:Fc Chimera

| | | | |
|---------------------------------|--|------------------|------------------------------|
| Catalog No. | CRH056A CRH056B CRH056C | Quantity: | 10 µg 3 x 10 µg 100 µg |
| Alternate Names: | NTN1L | | |
| Description: | <p>Netrin-1 controls guidance of CNS commissural axons and peripheral motor axons. Its association with either DCC or some UNC5 receptors will lead to axon attraction or repulsion, respectively. It also serve as a survival factor via its association with its receptors which prevent the initiation of apoptosis. Netrin-1 is also Involved in tumorigenesis by regulating apoptosis. Netrin-1 promotes atherosclerosis by retaining macrophages in the artery wall. Netrin-1 also governs induced pluripotent stem cell (iPS) formation and improves reprogramming efficiency of human and mouse somatic cells by limiting apoptosis mediated by Netrin-1 receptors DCC or UNC5b.</p> <p>Human Netrin-1 (aa 25-604) is fused at the C-terminus to the Fc portion of human IgG1.</p> | | |
| Gene ID: | 9423 | | |
| Protein Accession No: | O95631 | | |
| Source: | HEK293 cells | | |
| Molecular Weight: | ~90kDa (SDS-PAGE) | | |
| Formulation: | Lyophilized. Contains PBS plus protein stabilizer. No BSA included. The protein stabilizer shows no interference with protein activity. | | |
| Purity: | ≥95% (SDS-PAGE) | | |
| Endotoxin Level: | <0.1 EU/µg purified protein as determined by LAL test (Lonza). | | |
| Specificity: | Binds to human, mouse and rat UNC5B. | | |
| Biological Activity: | Induces pluripotent stem cell (iPS) formation and improves reprogramming efficiency of human and mouse somatic cells by inhibiting apoptosis mediated by the receptors DCC or UNC5B (at 150 ng/ml). Induces axon outgrowth. | | |
| Reconstitution: | Reconstitute 10 µg of protein with 100 µl (0.1 mg/ml) sterile water. PBS containing at least 0.1% BSA should be used for further dilutions. | | |
| Storage & Stability: | Store at 4°C upon arrival and at -20°C for long term. Lyophilized product is stable for at least 6 months after receipt when stored at -20°C. After reconstitution, prepare aliquots and store at -20°C. Avoid repeated freeze-thaw cycles. | | |



Recombinant Human Netrin-1:Fc triggers a DCC-dependent phosphorylation of ERK1/2.
Control HEK293 cells or netrin-1 receptor DCC expressing HEK293 cells were incubated with 5 nM Recombinant Human Netrin-1:Fc for 0 to 60 min. DCC, ERK1/2, and phospho-ERK1/3 were visualized by antibody chemiluminescence.



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
480 Neponset Street
Bldg 12A
Canton, MA 02021

Toll Free: 888-769-1246
Phone: 781-828-0610
Fax: 781-828-0542

E-mail: info@cellsciences.com
Website: www.cellsciences.com