


## PGF

### Recombinant Human Placenta Growth Factor, Animal Free

<b>Catalog No.</b>	CRH067A-AF CRH067B-AF CRH067C-AF	<b>Quantity:</b>	25 µg 100 µg 1.0 mg
<b>Alternate Names:</b>	PIGF		
<b>Description:</b>	<p>PIGF, a member of the VEGF family, was originally discovered in the placenta, where it was proposed to regulate vascular development and trophoblast growth and differentiation. Unlike VEGF, which is required for angiogenesis and endothelial cell maintenance and binds both Flt-1/sFlt-1 and Flk-1, PIGF is redundant for vascular development and selectively binds Flt-1/sFlt-1. PIGF stimulates endothelial cell growth, migration, and survival and plays a central role in pathologic angiogenesis, including in cancer and tissue ischemia. PIGF concentrations increase throughout pregnancy, peaking during the third trimester, and falling thereafter, probably as a consequence of placental maturation. Despite interest in the utility of PIGF for clinical diagnostics, the role of PIGF in preeclampsia and in placental development is not well understood.</p>		
<b>Gene ID:</b>	5228		
<b>UniProt ID:</b>	P49763		
<b>Source:</b>	<i>E. coli</i> <b>Manufactured without Animal-derived products, in an Animal Free facility.</b>		
<b>Molecular Weight:</b>	22.9/45.8 kDa Dimer, (204/408 aa)		
<b>Formulation:</b>	Lyophilized from a sterile filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA).		
<b>Purity:</b>	≥ 95% by reducing and non-reducing SDS-PAGE.		
<b>Endotoxin Level:</b>	≤1 EU/µg of protein by kinetic LAL analysis		
<b>Amino Acid Sequence:</b>	MLPAVPPQQW ALSAGNGSSE VEVVPFQEVW GRSYCRALER LVDVVSEYPS EVEHMFSPSC VSLLRCTGCC GDENLHCVPV ETANVTMQLL KIRSGDRPSY VELTFSQHVR CECRHSPGRQ SPDMPGDFRA DAPSFLLPPRR SLPMLFRMEW GCALTGSQSA VWPSSPVPEE IPRMHPGRNG KKQQRKPLRE KMKPERCGDA VPRR		
<b>Reconstitution:</b>	<b>Centrifuge vial prior to opening.</b> Add sterile distilled water to a concentration of 0.1 mg/ml. <b>DO NOT VORTEX.</b> Allow several minutes for complete reconstitution. Further dilutions should be made in appropriate buffered solutions.		
<b>Storage &amp; Stability:</b>	Store as supplied at -20°C to -80°C for up to 1 year. Upon reconstitution, prepare working aliquots and store at -20°C to -80°C. It is recommended that a carrier protein such as 0.1% HSA or BSA is added for long term storage. <b>Avoid repeated freeze-thaw cycles.</b>		





#### Human PlGF-3 Gel

Figure: 1 ug was run under non-reducing conditions (-) and reducing conditions (+) in a 4-20% Tris-Glycine gel, stained with Coomassie Blue. Human PlGF-3 is a dimer with a total predicted MW of 45.8 kDa (each monomer is 22.9 kDa).

**NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.**



**Cell Sciences®**  
65 Parker Street  
Unit 11  
Newburyport, MA 01950

Toll Free: 888-769-1246  
Phone: 978-572-1070  
Fax: 978-992-0298

E-mail: [info@cellsciences.com](mailto:info@cellsciences.com)  
Website: [www.cellsciences.com](http://www.cellsciences.com)