


BSG

Recombinant Human CD147 / EMMPRIN / Basigin (Fc Tag)

Catalog No.	CRH405A-Fc CRH405B-Fc	Quantity:	50 µg 100 µg
Alternate Names:	Basigin, 5F7, Collagenase stimulatory factor, Extracellular matrix metalloproteinase inducer, EMMPRIN, Leukocyte activation antigen M6, OK blood group antigen, Tumor cell-derived collagenase stimulatory factor, TCSF, CD147		
Description:	CD147/EMMPRIN (Extracellular Matrix Metalloproteinase Inducer), also known as Basigin (BSG), is a transmembrane glycoprotein with different forms resulted from different modes of glycosylation and N-terminal sequence variants. It is a member of the immunoglobulin superfamily with homology to both the immunoglobulin V domain and MHC class II antigen beta-chain. This protein play important roles in variety of events including spermatogenesis, embryo implantation, neural network formation. CD147 induces the production and release of matrix metalloproteinases (MMP) in the surrounding mesenchymal cells and tumor cells, and thereby promotes invasion, metastasis, growth and survival of malignant cells. Furthermore, CD147 also serves as a receptor for extracellular cyclophilinthe and its association with integrins might be important in signal transduction. Recently, CD147 displays increased expression in many cancers, and it has been previously demonstrated to participate in cancer metastasis and progression. Thus, CD147 and its antibody are used as an effective treatment for malignant cancers.		
UniProt ID:	P35613		
Accession Number:	NP_940991.1		
Protein Construction:	A DNA sequence encoding the extracellular domain (Met 1-His 205) of human CD147 (NP_940991.1) precursor was expressed with the fused Fc region of human IgG1 at the C-terminus.		
Source:	HEK293 Cells		
Formulation:	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.		
Molecular Weight:	The recombinant mature human CD147/Fc chimera protein consists of 422 amino acids and has a calculated molecular mass of 46.8 kDa. As a result of glycosylation, the recombinant protein migrates as an approximately 58-65 kDa band in SDS-PAGE under reducing conditions.		
Purity:	> 97 % as determined by SDS-PAGE.		
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method		
Biological Activity:	Testing in progress		
Predicted N-terminal:	Ala 22		

- Reconstitution:** **Centrifuge vial prior to opening.** Add sterile distilled water to a concentration of 0.1 mg/mL and gently pipette the solution up and down the sides of the vial. **DO NOT VORTEX.** Allow several minutes for complete reconstitution.
- Storage & Stability:** Stable for up to 1 year from date of receipt at -20°C to -80°C. After reconstitution, store working aliquots at -20°C to -80°C. **Avoid repeated freeze-thaw cycles.**

SDS-PAGE



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246
Phone: 978-572-1070
Fax: 978-992-0298

E-mail: info@cellsciences.com
Website: www.cellsciences.com