

## THPO

### Recombinant Human Thrombopoietin (Fc Tag)

<b>Catalog No.</b>	CRH470A-Fc CRH470B-Fc	<b>Quantity:</b>	20 µg 100 µg
<b>Alternate Names:</b>	Thrombopoietin, C-mpl ligand, ML, Megakaryocyte colony-stimulating factor, Megakaryocyte growth and development factor, MGDF, Myeloproliferative leukemia virus oncogene ligand		
<b>Description:</b>	Thrombopoietin also known as myeloproliferative leukemia virus ligand (c-Mpl), is a hematopoietic growth factor belonging to the EPO/TPO family. The thrombopoietin protein is produced mainly by the liver and the kidney that regulates the production of platelets by the bone marrow. Thrombopoietin protein stimulates both proliferation of progenitor megakaryocytes and their maturation to platelet-producing megakaryocytes, and also accelerates the recovery of platelets. Thrombopoietin protein is involved in cardiovascular disease as it regulates megakaryocyte development and enhances platelet adhesion/aggregation. It has been identified that surface c-MPL, the receptor for thrombopoietin protein, binds to the ligand and mediates the action.		
<b>UniProt ID:</b>	P40225		
<b>Accession Number:</b>	NP_000451.1		
<b>Protein Construction:</b>	A DNA sequence encoding the human THPO (Met1-Leu192) was expressed with the Fc region of human IgG1 at the C-terminus.		
<b>Source:</b>	HEK293 Cells		
<b>Formulation:</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.		
<b>Molecular Weight:</b>	The recombinant human THPO consists 409 amino acids and predicts a molecular mass of 45 kDa.		
<b>Purity:</b>	> 70 % as determined by SDS-PAGE		
<b>Endotoxin Level:</b>	< 1.0 EU per µg of the protein as determined by the LAL method.		
<b>Biological Activity:</b>	Testing in progress		
<b>Predicted N-terminal:</b>	Ser 22		
<b>Reconstitution:</b>	<b>Centrifuge vial prior to opening.</b> Add sterile distilled water to a concentration of 0.1 mg/mL and gently pipette the solution up and down the sides of the vial. <b>DO NOT VORTEX.</b> Allow several minutes for complete reconstitution.		
<b>Storage &amp; Stability:</b>	Stable for up to 1 year from date of receipt at -20°C to -80°C After reconstitution, store working aliquots at -20°C to -80°C. <b>Avoid repeated freeze-thaw cycles.</b>		



Cell Sciences®

65 Parker Street  
Unit 11  
Newburyport, MA 01950

Toll Free: 888-769-1246


Phone: 978-572-1070

Fax: 978-992-0298

E-mail: [info@cellsciences.com](mailto:info@cellsciences.com)

Website: [www.cellsciences.com](http://www.cellsciences.com)

SDS-PAGE



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



**Cell Sciences®**  
65 Parker Street  
Unit 11  
Newburyport, MA 01950

Toll Free: 888-769-1246  
Phone: 978-572-1070  
Fax: 978-992-0298

E-mail: [info@cellsciences.com](mailto:info@cellsciences.com)  
Website: [www.cellsciences.com](http://www.cellsciences.com)