

TNFRSF10A

Recombinant Human TRAIL-R1 / CD261 (Fc Tag)

Catalog No.	CRH474A-Fc CRH474B-Fc	Quantity:	100 µg 200 µg
Alternate Names:	Tumor necrosis factor receptor superfamily member 10A, Death receptor 4, TNF-related apoptosis-inducing ligand receptor 1, TRAIL receptor 1, TRAIL-R1, CD261		
Description:	Tumor necrosis factor receptor superfamily, member 1a (TRAIL R1) is a member of the TNF-receptor superfamily. This receptor is activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF1/TRAIL), and thus transduces cell death signal and induces cell apoptosis. Studies with FADD-deficient mice suggested that FADD, a death domain containing adaptor protein, is required for the apoptosis mediated by this protein. TRAIL R1 serves as a receptor for the cytotoxic ligand TNFSF1/TRAIL. The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. TRAIL R1 can promote the activation of NF-kappa-B. TRAIL R1 induces apoptosis of many transformed cell lines but not of normal tissues, even though its death domain-containing receptor, DR4, is expressed on both cell types.		
UniProt ID:	O00220		
Accession Number:	NP_003835		
Protein Construction:	A DNA sequence encoding the human TNFRSF10A extracellular domain (Met 1-Asn 239) was fused with the Fc region of human IgG1 at the C-terminus.		
Source:	HEK293 Cells		
Formulation:	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.		
Molecular Weight:	The recombinant human TNFRSF10A/Fc is a disulfide-linked homodimer. The reduced monomer consists of 372 amino acids and has a predicted molecular mass of 41.3 kDa. As a result of glycosylation, rh TNFRSF10A/Fc monomer migrates as an approximately 47 kDa band in SDS-PAGE under reducing conditions.		
Purity:	> 92 % as determined by SDS-PAGE		
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method.		
Biological Activity:	Testing in progress		
Predicted N-terminal:	Ala 109		
Reconstitution:	Centrifuge vial prior to opening. Add sterile distilled water to a concentration of 0.1 mg/mL and gently pipette the solution up and down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution.		
Storage & Stability:	Stable for up to 1 year from date of receipt at -20°C to -80°C After reconstitution, store working aliquots at -20°C to -80°C. Avoid repeated freeze-thaw cycles.		



cellsciences.com

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



® **Cell Sciences**®
65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246
Phone: 978-572-1070
Fax: 978-992-0298

E-mail: info@cellsciences.com
Website: www.cellsciences.com