

TNFRSF10B

Recombinant Human TRAIL-R2 / CD262 (Fc Tag)

Catalog No.	CRH490A-Fc	Quantity:	100 µg
	CRH490B-Fc		200 µg
	CRH490C-Fc		1.0 mg
Alternate Names:	Tumor necrosis factor receptor superfamily member 10B, Death receptor 5, TNF-related apoptosis-inducing ligand receptor 2, TRAIL receptor 2, TRAIL-R2, CD262		
Description:	TNF-related apoptosis-inducing ligand receptor 2 (TRAIL-R2), is a member of the TNF-receptor superfamily, and contains an intracellular death domain. This receptor can be activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF1/TRAIL/APO-2L), and transduces an apoptosis signal. Studies with FADD-deficient mice suggested that FADD, a death domain containing adaptor protein, is required for the apoptosis mediated by this protein. TRAIL-R2 was purified independently as the only receptor for TRAIL detectable on the surface of two different human cell lines that undergo apoptosis upon stimulation with TRAIL. TRAIL-R2 contains two extracellular cysteine-rich repeats, typical for TNF receptor (TNFR) family members, and a cytoplasmic death domain. TRAIL-R2 mediates apoptosis via the intracellular adaptor molecule FADD/MORT1. TRAIL receptors can signal both death and gene transcription, functions reminiscent of those of TNFR1 and TRAMP, two other members of the death receptor family. Defects in TRAIL-R2 may be a cause of head and neck squamous cell carcinomas (HNSCC) also known as squamous cell carcinoma of the head and neck.		
UniProt ID:	O14763		
Accession Number:	NP_003833		
Protein Construction:	A DNA sequence encoding the human TNFRSF10B (Met1-Glu182) was expressed with the Fc region of human IgG1 at the C-terminus.		
Source:	HEK293 Cells		
Formulation:	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.		
Molecular Weight:	The recombinant human TNFRSF10B consists of 365 amino acids and predicts a molecular mass of 41 kDa.		
Purity:	> 95 % as determined by SDS-PAGE		
Endotoxin Level:	< 1.0 EU per µg protein as determined by the LAL method.		
Biological Activity:	Testing in progress		
Predicted N-terminal:	Ile 56		
Reconstitution:	Centrifuge vial prior to opening. Add sterile distilled water to a concentration of 0.1 mg/mL and gently pipette the solution up and down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution.		



Cell Sciences®

65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246

Phone: 978-572-1070


Fax: 978-992-0298

E-mail: info@cellsciences.com

Website: www.cellsciences.com

Storage & Stability: Stable for up to 1 year from date of receipt at -20°C to -80°C
After reconstitution, store working aliquots at -20°C to -80°C.
Avoid repeated freeze-thaw cycles.

SDS-PAGE



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246
Phone: 978-572-1070
Fax: 978-992-0298

E-mail: info@cellsciences.com
Website: www.cellsciences.com