

FGFR4


Recombinant Human FGF Receptor 4 / CD334 (His & Fc Tag)

Catalog No.	CRH504A-HisFc CRH504B-HisFc	Quantity:	50 µg 100 µg
Alternate Names:	Fibroblast growth factor receptor 4, FGFR-4, CD334		
Description:	Fibroblast growth factor receptor 4 (CD334) is a member of the fibroblast growth factor receptor family, where amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein would consist of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of CD334 interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. CD334 preferentially binds acidic fibroblast growth factor and, although its specific function is unknown, it is overexpressed in gynecological tumor samples, suggesting a role in breast and ovarian tumorigenesis. CD334 signaling is down-regulated by receptor internalization and degradation; MMP14 promotes internalization and degradation of CD334. Mutations in CD334 lead to constitutive kinase activation or impair normal FGFR4 inactivation lead to aberrant signaling.		
UniProt ID:	P22455		
Accession Number:	NP_002002.3		
Protein Construction:	A DNA sequence encoding the extracellular domain (Met 1-Asp 369) of human FGFR4 precursor was expressed with the C-terminal polyhistidine tagged Fc region of human IgG1 at the C-terminus.		
Source:	HEK293 Cells		
Formulation:	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.		
Molecular Weight:	The recombinant human FGFR4/Fc is a disulfide-linked homodimer after removal of the signal peptide. The reduced monomer consists of 595 amino acids and has a predicted molecular mass of 66 kDa. In SDS-PAGE under reducing conditions, the rh FGFR4/Fc monomer migrates at ~100-110 kDa due to glycosylation.		
Purity:	> 97 % as determined by SDS-PAGE		
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method		
Biological Activity:	Measured by its ability to inhibit FGF-acidic dependent proliferation of Balb/c3T3 mouse fibroblasts. The ED50 for this effect is typically 10-60 ng/ml.		
Predicted N-terminal:	Leu 22		


Reconstitution: **Centrifuge vial prior to opening.** Add sterile distilled water to a concentration of 0.1 mg/mL and gently pipette the solution up and down the sides of the vial. **DO NOT VORTEX.** Allow several minutes for complete reconstitution.

Storage & Stability: Stable for up to 1 year from date of receipt at -20°C to -80°C. After reconstitution, store working aliquots at -20°C to -80°C. **Avoid repeated freeze-thaw cycles.**

Measured by its ability to inhibit FGF-acidic dependent proliferation of Balb/c3T3 mouse fibroblasts. The ED50 for this effect is typically 10-60 ng/ml.



SDS-PAGE



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.