

CEACAM5

Recombinant Human CEACAM5 / CD66e (Fc Tag)

Catalog No.	CRH614A-Fc	Quantity:	50 µg
	CRH614B-Fc		100 µg
Alternate Names:	Carcinoembryonic antigen-related cell adhesion molecule 5, Carcinoembryonic antigen, CEA, Meconium antigen 100, CD66e		
Description:	<p>CEACAM5 belongs to the large CEACAM subfamily of immunoglobulin superfamily. CEACAM5 is expressed primarily by epithelial cells. CEACAM5 contains one Ig-like V-type domain at the N-terminus, followed by six Ig-like C2-type domain and a GPI anchor, and exists as a homodimer. CEACAM5 and CEACAM6 are overexpressed in many cancers and are associated with adhesion and invasion. CEACAM5 can mediate cell-cell adhesion through homotypic and heterotypic interactions. It functions as a homotypic intercellular adhesion molecule and serves as a widely used tumor marker, since it is expressed at higher levels in tumorous tissues than in corresponding normal tissues. CEACAM5 has also been shown to contribute to tumorigenicity by inhibiting cellular differentiation. In addition, CEACAM5 is identified as the host receptor for the Dr family of adhesins of E.Coli, and the binding of E.coli Dr adhesins leads to dissociation of the CEACAM5 homodimer.</p>		
UniProt ID:	P06731		
Accession Number:	NP_000495.1		
Protein Construction:	A DNA sequence encoding the human CEACAM5 (Met1-Ala685) was expressed with the Fc region of human IgG1 at the C-terminus.		
Source:	HEK293 Cells		
Formulation:	<p>Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.</p>		
Molecular Weight:	The recombinant human CEACAM5 consists of 889 amino acids and predicts a molecular mass of 98.1 kDa.		
Purity:	> 95 % as determined by SDS-PAGE.		
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method		
Biological Activity:	Testing in progress		
Predicted N-terminal:	Lys 35		
Reconstitution:	<p>Centrifuge vial prior to opening. Add sterile distilled water to a concentration of 0.1 mg/mL and gently pipette the solution up and down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution.</p>		
Storage & Stability:	<p>Stable for up to 1 year from date of receipt at -20°C to -80°C After reconstitution, store working aliquots at -20°C to -80°C. Avoid repeated freeze-thaw cycles.</p>		



Cell Sciences®

65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246


Phone: 978-572-1070

Fax: 978-992-0298

E-mail: info@cellsciences.com

Website: www.cellsciences.com

SDS-PAGE



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246
Phone: 978-572-1070
Fax: 978-992-0298

E-mail: info@cellsciences.com
Website: www.cellsciences.com