


## CD22

### Recombinant Human Siglec-2 / CD22 (aa 176-687, His Tag)

<b>Catalog No.</b>	CRH658A-His CRH658B-His	<b>Quantity:</b>	50 µg 100 µg
<b>Alternate Names:</b>	B-cell receptor CD22, B-lymphocyte cell adhesion molecule, BL-CAM, Sialic acid-binding Ig-like lectin 2, Siglec-2, T-cell surface antigen Leu-14, CD22		
<b>Description:</b>	CD22 is a member of the immunoglobulin superfamily, SIGLEC family of lectins. It is first expressed in the cytoplasm of pro-B and pre-B cells, and on the surface as B cells mature to become IgD+. CD22 serves as an adhesion receptor for sialic acid-bearing ligands expressed on erythrocytes and all leukocyte classes. In addition to its potential role as a mediator of intercellular interactions, signal transduction through CD22 can activate B cells and modulate antigen receptor signaling in vitro. The phenotype of CD22-deficient mice suggests that CD22 is primarily involved in the generation of mature B cells within the bone marrow, blood, and marginal zones of lymphoid tissues. CD22 recruits the tyrosine phosphatase Src homology 2 domain-containing phosphatase 1 (SHP-1) to immunoreceptor tyrosine-based inhibitory motifs (ITIMs) and inhibits B-cell receptor (BCR)-induced Ca <sup>2+</sup> signaling on normal B cells. CD22 interacts specifically with ligands carrying alpha2-6-linked sialic acids. As an inhibitory coreceptor of the B-cell receptor (BCR), CD22 plays a critical role in establishing signaling thresholds for B-cell activation.		
<b>UniProt ID:</b>	P20273		
<b>Accession Number:</b>	NP_001762.2		
<b>Protein Construction:</b>	A DNA sequence encoding the human CD22 (Trp176-Arg687) was fused with a polyhistidine tag at the C-terminus.		
<b>Source:</b>	HEK293 Cells		
<b>Formulation:</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.		
<b>Molecular Weight:</b>	The rhCD22 consists of 523 aa with a predicted MW of 58.5 kDa. In SDS-PAGE under reducing conditions, rhCD22 migrates at ~90-100 kDa due to glycosylation.		
<b>Purity:</b>	> 90 % as determined by SDS-PAGE.		
<b>Endotoxin Level:</b>	< 1.0 EU per µg of the protein as determined by the LAL method		
<b>Biological Activity:</b>	Testing in progress		
<b>Predicted N-terminal:</b>	Trp 176		
<b>Reconstitution:</b>	<b>Centrifuge vial prior to opening.</b> Add sterile distilled water to a concentration of 0.1 mg/mL and gently pipette the solution up and down the sides of the vial. <b>DO NOT VORTEX.</b> Allow several minutes for complete reconstitution.		
<b>Storage &amp; Stability:</b>	Stable for up to 1 year from date of receipt at -20°C to -80°C After reconstitution, store working aliquots at -20°C to -80°C. <b>Avoid repeated freeze-thaw cycles.</b>		



SDS-PAGE



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



**Cell Sciences®**  
65 Parker Street  
Unit 11  
Newburyport, MA 01950

Toll Free: 888-769-1246  
Phone: 978-572-1070  
Fax: 978-992-0298

E-mail: [info@cellsciences.com](mailto:info@cellsciences.com)  
Website: [www.cellsciences.com](http://www.cellsciences.com)