

NGFR

Recombinant Human Nerve Growth Factor Receptor / CD271 (Fc Tag)

Catalog No.	CRH698A-Fc CRH698B-Fc	Quantity:	50 µg 100 µg
Alternate Names:	Tumor necrosis factor receptor superfamily member 16, Gp80-LNGFR, Low affinity neurotrophin receptor p75NTR, Low-affinity nerve growth factor receptor, NGF receptor, p75 ICD, CD271		
Description:	Nerve growth factor receptors belong to a large growth factor receptor family made of two types of receptors: high-affinity nerve growth factor receptor and low-affinity nerve growth factor receptor. High-affinity nerve growth factor receptor is also referred as Trk family whose members are bound by some neurotrophins with high affinity. The Low-affinity nerve growth factor receptor also named p75 which binds with all kinds of neurotrophins with low affinity. All the four kinds of neurotrophins, including Nerve growth factor, Brain derived neurotrophic factor, Neurotrophin-3, and Neurotrophin-4 bind to the p75. Studies have proved that NGFR acts as a molecular signal switch that determines cell death or survival by three steps. First, pro-nerve growth factor (prNGF) triggers cell apoptosis by its high affinity binding to p75NTR, while NGF induces neuronal survival with low-affinity binding. Second, p75NTR mediates cell death by combining with co-receptor sortilin, whereas it promotes neuronal survival through combination with proNGF. Third, release of the intracellular domain chopper or cleavage short p75 NTR can independently initiate neuronal apoptosis.		
UniProt ID:	P08138		
Protein Construction:	A DNA sequence encoding the human NGFR extracellular domain (Met 1-Asn 250) was fused with the Fc region of human IgG1 at the C-terminus.		
Source:	HEK293 Cells		
Formulation:	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.		
Molecular Weight:	The secreted rhNGFR/Fc is a disulfide-linked homodimer. The reduced monomer consists of 463 aa with a predicted MW of 50.6 kDa and migrates at ~70-80 kDa in SDS-PAGE under reducing conditions due to glycosylation.		
Purity:	> 95 % as determined by SDS-PAGE.		
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method		
Biological Activity:	Measured by its ability to inhibit NGF-dependent proliferation of TF-1 human erythroleukemic cells. The ED50 for this effect is typically 0.5-3 µg/mL in the presence of 4 ng/mL Recombinant Human NGF.		
Predicted N-terminal:	Lys 29		



Reconstitution:

Centrifuge vial prior to opening. Add sterile distilled water to a concentration of 0.1 mg/mL and gently pipette the solution up and down the sides of the vial.

DO NOT VORTEX. Allow several minutes for complete reconstitution.


Storage & Stability:

Stable for up to 1 year from date of receipt at -20°C to -80°C

After reconstitution, store working aliquots at -20°C to -80°C.

Avoid repeated freeze-thaw cycles.

SDS-PAGE



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246
Phone: 978-572-1070
Fax: 978-992-0298

E-mail: info@cellsciences.com
Website: www.cellsciences.com