cellsciences.com

PCSK9

Recombinant Human Proprotein Convertase 9 / NARC-1 (D374Y) (Fc Tag)

Catalog No. CRH755A-Fc **Quantity**: 10 μg

CRH755B-Fc 50 μg

Alternate Names: Proprotein convertase subtilisin/kexin type 9, Neural apoptosis-regulated convertase 1,

NARC-1, Proprotein convertase 9, PC9, Subtilisin/kexin-like protease PC9

Description: Proprotein convertase 9 (PC9) is a newly identified human secretory subtilase belonging

to the proteinase K subfamily of the secretory subtilase family. PC9 is an enzyme which in humans is encoded by the PCSK9 gene with orthologs found across many species. It is expressed in neuroepithelioma, colon carcinoma, hepatic and pancreatic cell lines, and in Schwann cells. PC9 protein is highly expressed in the liver and regulates low density lipoprotein receptor (LDLR) protein levels. Inhibition of PC9 protein function is currently

being explored as a means of lowering cholesterol levels.

UniProt ID: Q8NBP7

Accession Number: NP_777596.2

Protein Construction: A DNA sequence encoding the human PCSK9 (Met1-Gln692), with mutation Asp 374

Tyr, was expressed with the Fc region of mouse IgG1 at the C-terminus.

Source: HEK293 Cells

Formulation: Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants

before lyophilization.

Molecular Weight: The recombinant human PCSK9 consists of 896 amino acids and predicts a molecular

mass of 97.5 kDa.

Purity: > 85 % as determined by SDS-PAGE.

Endotoxin Level: $< 1.0 \text{ EU per } \mu\text{g}$ protein as determined by the LAL method.

Biological Activity: Testing in progress

Predicted N-terminal: Gln 31

Reconstitution: Centrifuge vial prior to opening. Add sterile distilled water to a concentration of 0.1

mg/mL and gently pipette the solution up and down the sides of the vial. **DO NOT VORTEX**. Allow several minutes for complete reconstitution.

E-mail: info@cellsciences.com

Website: www.cellsciences.com

Storage & Stability: Stable for up to 1 year from date of receipt at -20°C to -80°C


After reconstitution, store working aliquots at -20°C to -80°C.

Toll Free: 888-769-1246

Phone: 978-572-1070


Fax: 978-992-0298

Avoid repeated freeze-thaw cycles.



cellsciences.com

SDS-PAGE



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

Toll Free: 888-769-1246 E-mail: Phone: 978-572-1070 Website

Fax: 978-992-0298

E-mail: <u>info@cellsciences.com</u>
Website: <u>www.cellsciences.com</u>