cellsciences.com

IL₂

Recombinant Mouse Interleukin-2, Animal Free

Catalog No. CRI145A-AF **Quantity**: 5 μg

CRI145B-AF 20 μg CRI145C-AF 1.0 mg

Alternate Names: IL-2, lymphokine, aldesleukin, T cell growth factor

Description: Interleukin-2 (IL-2) is an immunomodulatory cytokine that is produced by lymphocytes. IL

-2 signals through the IL-2R receptor to induce activated T cell proliferation and promote T cell differentiation. IL-2 also stimulates the proliferation and differentiation of B cells, natural killer cells, monocytes, and macrophages. IL-2 plays a central role in the expansion and maintenance of regulatory T cells, although it inhibits the development of

Th17 polarized cells.

Gene ID: 16183

UniProt ID: P04351

Source: E. coli

Made without animal-derived materials in an animal free facility.

Molecular Weight: Monomer, 17.4 kDa (150 aa)

Formulation: Lyophilized from sterile filtered aqueous solution containing 10 mM sodium citrate, pH

4.0.

Purity: >95% by nonreducing and reducing SDS-PAGE

Endotoxin Level: $\leq 1 \text{ EU/µg}$

Biological Activity: ED₅₀ \leq 5 ng/ml, as determined by proliferation assay of murine CTLL-2 cells.

Specific Activity: $\geq 2.0 \times 10^5 \text{ U/mg}$ calibrated against recombinant mouse IL-2 WHO International

Standard (NIBSC code: 93/566).

Amino Acid Sequence: MAPTSSSTSS STAEAQQQQQ QQQQQQQHLE QLLMDLQELL SRMENYRNLK

LPRMLTFKFY LPKQATELKD LQCLEDELGP LRHVLDLTQS KSFQLEDAEN FISNIRVTVV KLKGSDNTFE CQFDDESATV VDFLRRWIAF CQSIISTSPQ

Reconstitution: Centrifuge vial prior to opening. Add sterile 10mM acetic acid to a concentration of 0.1

mg/mL. DO NOT VORTEX. Allow several minutes for complete reconstitution. Further

dilutions should be made in appropriate buffered solutions.

Toll Free: 888-769-1246

Phone: 978-572-1070

Fax: 978-992-0298

Storage & Stability: Store as supplied at -20 °C to -80 °C for up to 1 year. Upon reconstitution, the

preparation is stable for up to one month at 2-8°C. For long term storage, freeze in working aliquots containing 0.1% BSA at -20 °C to -80 °C. **Avoid repeated freeze-thaw**


E-mail: info@cellsciences.com


Website: www.cellsciences.com

cycles.

cellsciences.com

Figure: 1 µg run under (+) reducing conditions in a 4 -20% Tris-Glycine gel, stained with Coomassie Blue.





NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

Toll Free: 888-769-1246 Phone: 978-572-1070

Fax: 978-992-0298

E-mail: info@cellsciences.com www.cellsciences.com Website: