## cellsciences.com

#### **II10**

### **Recombinant Rat Interleukin-10**

**Catalog No.** CRI155A **Quantity:** 2 μg

CRI155B 10 μg CRI155C 1.0 mg

Alternate Names: IL-10, Cytokine synthesis inhibitory factor, CSIF

**Description:** IL10 is a cytokine produced primarily by monocytes and to a lesser extent by

lymphocytes. This cytokine has pleiotropic effects in immunoregulation and inflammation. It down-regulates the expression of Th1 cytokines, MHC class II Ags, and costimulatory molecules on macrophages. It also enhances B cell survival, proliferation, and antibody production. This cytokine can block NF-kappa B activity, and is involved in the regulation of the JAK-STAT signaling pathway. Knockout studies in mice suggest the function of this

cytokine as an essential immunoregulator in the intestinal tract.

UniProt ID: P29456

**Gene ID:** 25325

Source: E. coli

Molecular Weight: 18.8 kDa (161 aa) monomer

Formulation: Lyophilized from a sterile filtered aqueous solution containing 0.1% Trifluoroacetic Acid

(TFA)

**Purity:**  $\geq$  95% by reducing and non-reducing SDS-PAGE

**Endotoxin Level:** < 1 EU/µg by kinetic LAL analysis.

Amino Acid Sequence: MSKGHSIRGD NNCTHFPVSQ THMLRELRAA FSQVKTFFQK KDQLDNILLT

DSLLQDFKGY LGCQALSEMI KFYLVEVMPQ AENHGPEIKE HLNSLGEKLK TLWIQLRRCH RFLPCENKSK AVEQVKNDFN KLQDKGVYKA MNEFDIFINC

**IEAYVTLKMK N** 

**Reconstitution:** Centrifuge vial prior to opening. Add sterile distilled water to a concentration of 0.1

mg/mL and gently pipet the solution up and down the sides of the vial.

DO NOT VORTEX. Allow several minutes for reconstitution. A small amount of

precipitate may be seen.

**Storage & Stability:** Store as supplied at -20°C to -80°C for up to 1 year. Upon reconstitution, prepare

working aliquots and store at -20°C to -80°C. It is recommended that a carrier protein

E-mail: info@cellsciences.com

Website: www.cellsciences.com

such as 0.1% HSA or BSA is added for long term storage.


Toll Free: 888-769-1246

Phone: 978-572-1070

Fax: 978-992-0298

Avoid repeated freeze-thaw cycles.

# cellsciences.com



#### Rat IL-10 Gel

Figure: 1 ug run under (-) non-reducing and (+) reducing conditions in a 4-20% Tris-Glycine gel, stained with Coomassie Blue. Rat IL-10 is predicted to have a

MW of 18.8 kDa.

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

Toll Free: 888-769-1246 Phone: 978-572-1070 Fax: 978-992-0298

E-mail: info@cellsciences.com Website: www.cellsciences.com