



Gdnf

Recombinant Mouse Glial Derived Neurotrophic Factor-Animal Free

Catalog No.	CRM014A-AF CRM014B-AF CRM014C-AF CRM014D-AF	Quantity:	2 µg 10 µg 1 mg 100 µg
Alternate Names:	GDNF, astrocyte-derived trophic factor, ATF1, ATF2, HSCR3, HFB1-GDNF		
Description:	Glial Cell Line-Derived Neurotrophic Factor (GDNF) is a neurotrophic factor that is closely related to other neurotrophic factors, such as Neurturin, Persephin, and Artemin, by a common structural feature called the cysteine-knot. GDNF signals through a multicomponent system of receptors that includes RET and GFRα1-4, to promote dopamine uptake, survival and differentiation of neurons.		
Physical Appearance:	Sterile filtered white lyophilized (freeze-dried) powder.		
Gene ID:	2668		
UniProt ID:	P48540		
Source:	<i>E. coli</i>		
Molecular Weight:	Dimer, 15.1/30.2 kDa (135/270 aa)		
Formulation:	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA).		
Purity:	≥95% as determined by reducing and non-reducing SDS-PAGE		
Endotoxin Level:	≤ 0.1 EU/µg, by kinetic LAL		
Amino Acid Sequence:	MSPDKQAAL PRRERNRQAA AASPENSRGK GRRGQRGKNR GCVLTAIHLN VTDLGLGYET KEELIFRYCS GSCESAETMY DKILKNLSRS RRLTSDKVGQ ACCRPVAFFDD DLSFLDDNLV YHILRKHS AK RCGCI		
Reconstitution:	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. DO NOT VORTEX . It is recommended to reconstitute the lyophilized product with sterile water at a concentration of 0.1 mg/ml, which can be further diluted into other aqueous solutions.		
Storage & Stability:	Store as supplied at -20°C to -80°C. Upon reconstitution, prepare working aliquots and store at -20°C to -80°C. It is recommended to add a carrier protein such as 0.1% HSA or BSA for long term storage. This depends upon the particular application employed. Avoid repeated freeze-thaw cycles.		



Figure 1: C6 cells were cultured with 0 to 5 $\mu\text{g/ml}$ Mouse GDNF. Cell proliferation was measured after 7 days and the linear portion of the curve was used to calculate the ED50.



Mouse GDNF Gel

Figure: 1 μg run under (-) non-reducing conditions and (+) reducing conditions in a 4-20% Tris-Glycine gel, stained with Coomassie Blue. Mouse GDNF is a dimer with a predicted MW of 30.2 kDa (each monomer is 15.1 kDa).

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246
Phone: 978-572-1070
Fax: 978-992-0298

E-mail: info@cellsciences.com
Website: www.cellsciences.com