

## Pdcd2lg2

### Recombinant Mouse PD-L2/CD273:Fc Chimera

<b>Catalog No.</b>	CRM037	<b>Quantity:</b>	100 µg
<b>Alternate Names:</b>	Programmed Cell Death 1 Ligand 2, B7-DC, Btdc, PD-L2, PDL2, CD273		
<b>Description:</b>	<p>T cells require a signal induced by the engagement of the T cell receptor and a costimulatory signal(s) through distinct T cell surface molecules for optimal T cell activation and tolerance. CD273 (PD-L2) is one of two ligands for programmed death-1 (PD-1; CD279), a member of the CD28 family of immunoreceptors. The other identified ligand is PD-L1. CD273 is broadly expressed and also up regulated in a variety of tumor cell lines. On previously activated T cells, CD273 interaction with PD-1 inhibits TCR mediated proliferation and cytokine production, suggesting an inhibitory role in regulating immune responses. CD273 has a costimulatory function on resting T cells activated with suboptimal TCR signals.</p> <p>The extracellular domain of mouse CD273 [PD-L2] (aa 20-219) is fused to the N-terminus of the Fc region of mouse IgG2a.</p>		
<b>Gene ID:</b>	58205		
<b>Protein Accession No:</b>	NP_067371.1		
<b>Source:</b>	CHO cells		
<b>Formulation:</b>	Lyophilized from a 0.2 µm filtered solution containing PBS.		
<b>Purity:</b>	≥98% (SDS-PAGE)		
<b>Endotoxin Level:</b>	<0.06 EU/µg purified protein as determined by LAL test (Lonza).		
<b>Reconstitution:</b>	Reconstitute at 100 µg/ml in sterile PBS.		
<b>Storage &amp; Stability:</b>	Store at 4°C upon arrival and at -20°C for long term. Lyophilized product is stable for at least 1 year after receipt when stored at -20°C. After reconstitution, prepare aliquots and store at -20°C. Stable for up to 3 month at -20°C. <b>Avoid repeated freeze-thaw cycles.</b>		

**NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.**

