

Ghr


Recombinant Mouse Growth Hormone Receptor (His Tag)

Catalog No.	CRM517A-His CRM517B-His	Quantity:	50 µg 100 µg
Alternate Names:	Growth hormone receptor, GH receptor, Somatotropin receptor, Growth hormone-binding protein, GH-binding protein, GHBP, Serum-binding protein		
Description:	Growth hormone receptor (GHR) is a single-pass type I membrane protein containing one fibronectin type-III domain. GHR is expressed in various tissues with high expression in liver and skeletal muscle. Isoform 4 of GHR is predominantly expressed in kidney, bladder, adrenal gland, placental villi and brain stem. Isoform 1 expression of GHR in the placenta is predominant in chorion and decidua. Isoform 2 of GHR is expressed in lung, stomach and muscle. GHR is a receptor for pituitary gland growth hormone. It is involved in regulating postnatal body growth. On ligand binding, it couples to the JAK2 / STAT5 pathway. Isoform 2 of GHR up-regulates the production of GHBP and acts as a negative inhibitor of GH signaling. Defects in GHR are a cause of Laron syndrome which is a severe form of growth hormone insensitivity characterized by growth impairment, short stature, dysfunctional growth hormone receptor, and failure to generate insulin-like growth factor I in response to growth hormone. Defects in GHR may also be a cause of idiopathic short stature autosomal (ISSA) which is defined by a subnormal rate of growth.		
UniProt ID:	P16882		
Accession Number:	NP_034414.2		
Protein Construction:	A DNA sequence encoding the extracellular domain of mouse GHR (Met 1-Gln 273) was fused with a polyhistidine tag at the C-terminus.		
Source:	HEK293 Cells		
Formulation:	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.		
Molecular Weight:	The rhGHR consists of 260 aa with a predicted MW of 30.3 kDa and migrates at ~40-45 kDa in SDS-PAGE under reducing conditions, due to glycosylation.		
Purity:	> 95 % as determined by SDS-PAGE.		
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method		
Biological Activity:	Measured by its ability to inhibit proliferation of INS-1 cells induced by human growth hormone. The ED50 for this effect is 2-8µg/mL in the presence of 50 ng/mL human growth hormone.		
Predicted N-terminal:	Thr 25		
Reconstitution:	Centrifuge vial prior to opening. Add sterile distilled water to a concentration of 0.1 mg/mL and gently pipette the solution up and down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution.		



Storage & Stability: Stable for up to 1 year from date of receipt at -20°C to -80°C
After reconstitution, store working aliquots at -20°C to -80°C.
Avoid repeated freeze-thaw cycles.

Measured by its ability to inhibit proliferation of INS-1 cells induced by human growth hormone. The ED50 for this effect is 2-8µg/mL in the presence of 50 ng/mL human growth hormone.



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246
Phone: 978-572-1070
Fax: 978-992-0298

E-mail: info@cellsciences.com
Website: www.cellsciences.com