

Icoslg

Recombinant Mouse ICOS Ligand / B7-H2 / CD275 (His & Fc Tag)


Catalog No.	CRM567A-HisFc CRM567B-HisFc	Quantity:	100 µg 200 µg
Alternate Names:	ICOS ligand, B7 homolog 2, B7-H2, B7-like protein GI50, B7-related protein 1, B7RP-1, LICOS, CD275		
Description:	Inducible co-stimulator ligand (ICOSL), also known as B7-H2, is a member of the B7 family of co-stimulatory molecules related to B7-1 and B7-2. It is a transmembrane glycoprotein with extracellular IgV and IgC domains, and binds to ICOS on activated T cells, thus delivers a positive costimulatory signal for optimal T cell function. The structural features of ICOSL are crucial for its costimulatory function. Present study shows that ICOSL displays a marked oligomerization potential, resembling more like B7-1 than B7-2. B7-H2-dependent signaling may play an active role in a proliferative response rather than in cytokine and chemokine production. The CD28/B7 and ICOS/B7-H2 pathways are both critical for costimulating T cell immune responses. Deficiency in either pathway results in defective T cell activation, cytokine production and germinal center formation.		
UniProt ID:	Q9JHJ8		
Accession Number:	NP_056605.1		
Protein Construction:	A DNA sequence encoding the mouse B7-H2 extracellular domain (Met 1-Lys 279) was fused with the C-terminal polyhistidine-tagged Fc region of human IgG1 at the C-terminus.		
Source:	HEK293 Cells		
Formulation:	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.		
Molecular Weight:	The secreted rmB7-H2/Fc is a disulfide-linked homodimer after removal of the signal peptide. The reduced monomer consists of 481 aa with a predicted MW of 54.3 kDa and migrates at ~75-85 kDa in reduced SDS-PAGE, due to glycosylation.		
Purity:	> 95 % as determined by SDS-PAGE.		
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method		
Biological Activity:	In a functional ELISA, immobilized human ICOS at 1 µg/ml (100 µl/well) can bind biotinylated mouse B7-H2 Fc chimera with a linear range of 0.125-1.0 µg/ml.		
Predicted N-terminal:	Glu 47		
Reconstitution:	Centrifuge vial prior to opening. Add sterile distilled water to a concentration of 0.1 mg/mL and gently pipette the solution up and down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution.		



Storage & Stability:

Stable for up to 1 year from date of receipt at -20°C to -80°C
After reconstitution, store working aliquots at -20°C to -80°C.
Avoid repeated freeze-thaw cycles.

SDS-PAGE



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246
Phone: 978-572-1070
Fax: 978-992-0298

E-mail: info@cellsciences.com
Website: www.cellsciences.com