


Bsg

Recombinant Mouse CD147 / Basigin (His Tag)

Catalog No.	CRM590A-His CRM590B-His	Quantity:	50 µg 100 µg
Alternate Names:	Basigin, Basic immunoglobulin superfamily, HT7 antigen, Membrane glycoprotein gp42, CD147		
Description:	Basigin (CD147) is a transmembrane glycoprotein with different forms resulting from different modes of glycosylation and N-terminal sequence variants. It is a member of the immunoglobulin superfamily with homology to both the immunoglobulin V domain and MHC class II antigen beta-chain. This protein play important roles in variety of events including spermatogenesis, embryo implantation, neural network formation. CD147 induces the production and release of matrix metalloproteinases (MMP) in the surrounding mesenchymal cells and tumor cells, and thereby promotes invasion, metastasis, growth and survival of malignant cells. Furthermore, CD147 also serves as a receptor for extracellular cyclophilinthe and its association with integrins might be important in signal transduction. Recently, CD147 displays increased expression in many cancers, and it has been previously demonstrated to participate in cancer metastasis and progression. Thus, CD147 and its antibody are used as an effective treatment for malignant cancers.		
UniProt ID:	P18572-2		
Accession Number:	NP_001070652.1		
Protein Construction:	A DNA sequence encoding the extracellular domain of mouse BSG isoform 2 (Met 1-Arg 209) was expressed, fused with a polyhistidine tag at the C-terminus.		
Source:	HEK293 Cells		
Formulation:	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.		
Molecular Weight:	The secreted rmCD147 consists of 199 aa with a predicted MW of 30 kDa and migrates at ~30-35 kDa in SDS-PAGE under reducing conditions, due to glycosylation.		
Purity:	> 90 % as determined by SDS-PAGE.		
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method		
Biological Activity:	Testing in progress		
Predicted N-terminal:	Ala 22		
Reconstitution:	Centrifuge vial prior to opening. Add sterile distilled water to a concentration of 0.1 mg/mL and gently pipette the solution up and down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution.		
Storage & Stability:	Stable for up to 1 year from date of receipt at -20°C to -80°C After reconstitution, store working aliquots at -20°C to -80°C. Avoid repeated freeze-thaw cycles.		

SDS-PAGE



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246
Phone: 978-572-1070
Fax: 978-992-0298

E-mail: info@cellsciences.com
Website: www.cellsciences.com