


Cd99I2

Recombinant Mouse Ephrin-A1 / CD99 (His Tag)

Catalog No.	CRM656A-His CRM656B-His	Quantity:	50 µg 100 µg
Alternate Names:	CD99 antigen-like protein 2, MIC2-like protein 1, CD99		
Description:	EPH-related receptor tyrosine kinase ligand 1 (Ephrin-A1) is a member of the ephrin (EPH) family. The Eph family receptor interacting proteins (ephrins) are a family of proteins that serve as the ligands of the Eph receptor, which compose the largest known subfamily of receptor protein-tyrosine kinases (RTKs). Ephrin-A1 and its Eph family of receptor tyrosine kinases are expressed by cells of the SVZ. Ephrin subclasses are further distinguished by their mode of attachment to the plasma membrane: ephrin-A ligands bind EphA receptors and are anchored to the plasma membrane via a glycosylphosphatidylinositol (GPI) linkage, whereas ephrin-B ligands bind EphB receptors and are anchored via a transmembrane domain. An exception is the EphA4 receptor, which binds both subclasses of ephrins. Ephrin-A1 and one of its receptor EphA2 were expressed in xenograft endothelial cells and also tumor cells and play a role in human cancers, at least in part by influencing tumor neovascularization.		
UniProt ID:	Q8BIF0		
Accession Number:	NP_612182.1		
Protein Construction:	A DNA sequence encoding the mouse EphA1 without the pro peptide (Met 1-Ser 182) was fused with a polyhistidine tag at the C-terminus.		
Source:	HEK293 Cells		
Formulation:	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.		
Molecular Weight:	The secreted rmEphA1 consists of 175 aa with a predicted MW of 20 kDa and migrates at ~27 kDa in SDS-PAGE under reducing conditions, due to glycosylation.		
Purity:	> 97 % as determined by SDS-PAGE.		
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method		
Biological Activity:	Measured by its ability to compete with mouse EphA1 for binding to immobilized mouse EphA2 in a functional ELISA assay.		
Predicted N-terminal:	Asp 19		
Reconstitution:	Centrifuge vial prior to opening. Add sterile distilled water to a concentration of 0.1 mg/mL and gently pipette the solution up and down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution.		
Storage & Stability:	Stable for up to 1 year from date of receipt at -20°C to -80°C After reconstitution, store working aliquots at -20°C to -80°C. Avoid repeated freeze-thaw cycles.		



SDS-PAGE



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.