

## Ifngr2


### Recombinant Mouse IFN-gamma Receptor 2 (His Tag)

<b>Catalog No.</b>	CRM674A-His CRM674B-His	<b>Quantity:</b>	50 µg 100 µg
<b>Alternate Names:</b>	Ifngr2 protein, Interferon gamma receptor 2		
<b>Description:</b>	Interferon gamma receptor beta chain (IFNGR2) belongs to the type II cytokine receptor family whose deficiency is a cause of autosomal recessive mendelian susceptibility to mycobacterial disease (MSMD), also known as familial disseminated atypical mycobacterial infection. This accessory factor is an integral part of the IFN-gamma signal transduction pathway and is likely to interact with GAF, JAK1, and/or JAK2. IFNGR2 is a component of the IFNgamma receptor complex along with the IFNgammaR alpha chain (IFNGR1), and is a new Bax suppressor. The C-terminal fragment (cytoplasmic domain) of IFNGR2 is expressed in human cancer cell lines of megakaryocytic cancer (DAMI), breast cancer (MDA-MD-468), and prostate cancer (PC3 cells). The Th1 cytokine IFNgamma, acting through its heterodimeric receptors, IFNGR1 and IFNGR2, in the induction/proliferation of Th1 cells, might suppress the Th2 responses that may underlie atopic asthma. IFNGR2 has always been seen as a key mechanism for shielding T lymphocytes from the antiproliferative effects of the IFNgamma-signal transducer and activator of transcription 1 (STAT1) pathway.		
<b>UniProt ID:</b>	Q63953		
<b>Accession Number:</b>	NP_032364.1		
<b>Protein Construction:</b>	A DNA sequence encoding the extracellular domain of mouse IFNGR2 (Met 1-Val 243) was expressed, with a C-terminal polyhistidine tag.		
<b>Source:</b>	HEK293 Cells		
<b>Formulation:</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.		
<b>Molecular Weight:</b>	The secreted rmIFNGR2 consists of 234 aa with a predicted MW of 26.7 kDa and migrates at ~40-45 kDa in SDS-PAGE under reducing conditions, due to glycosylation.		
<b>Purity:</b>	> 97 % as determined by SDS-PAGE.		
<b>Endotoxin Level:</b>	< 1.0 EU per µg of the protein as determined by the LAL method		
<b>Biological Activity:</b>	Testing in progress		
<b>Predicted N-terminal:</b>	Ser 20		
<b>Reconstitution:</b>	<b>Centrifuge vial prior to opening.</b> Add sterile distilled water to a concentration of 0.1 mg/mL and gently pipette the solution up and down the sides of the vial. <b>DO NOT VORTEX.</b> Allow several minutes for complete reconstitution.		



**Storage & Stability:** Stable for up to 1 year from date of receipt at -20°C to -80°C  
After reconstitution, store working aliquots at -20°C to -80°C.  
**Avoid repeated freeze-thaw cycles.**

SDS-PAGE



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®  
65 Parker Street  
Unit 11  
Newburyport, MA 01950

Toll Free: 888-769-1246  
Phone: 978-572-1070  
Fax: 978-992-0298

E-mail: [info@cellsciences.com](mailto:info@cellsciences.com)  
Website: [www.cellsciences.com](http://www.cellsciences.com)