


Dpp4

Recombinant Mouse DPP4 / CD26 (Fc Tag)

Catalog No.	CRM688A-Fc CRM688B-Fc	Quantity:	10 µg 20 µg
Alternate Names:	Dipeptidyl peptidase 4, Dipeptidyl peptidase IV, DPP IV, T-cell activation antigen CD26, Thymocyte-activating molecule, THAM, CD26		
Description:	Dipeptidyl peptidase-4 (DPP4) is a serine exopeptidase belonging to the S9B protein family that cleaves X-proline dipeptides from the N-terminus of proteins. The enzyme is a type II transmembrane glycoprotein, expressed on the surface of many cell types. It is also present in serum and other body fluids in a truncated form (sCD26/DPPIV). The soluble CD26 (sCD26) is a tumor marker for the detection of colorectal cancer (CRC) and advanced adenomas. As both a regulatory enzyme and a signalling factor, DPP4 has been evaluated and described in many studies. DPP4 inhibition results in increased blood concentration of the incretin hormones glucagon-like peptide-1 (GLP-1) and gastric inhibitory polypeptide (GIP). This causes an increase in glucose-dependent stimulation, resulting in a lowering of blood glucose levels. Recent studies have shown that DPP4 inhibitors can induce a significant reduction in glycosylated hemoglobin HbA(1c) levels, alone or in combination with other antidiabetic agents, and are a new class of drugs for treating Type 2 diabetes.		
UniProt ID:	P28843		
Accession Number:	NP_034204.1		
Protein Construction:	A DNA sequence encoding the mouse Dpp4 (Ser29-His760) was expressed with the Fc region of human IgG1 at the N-terminus.		
Source:	HEK293 Cells		
Formulation:	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.		
Molecular Weight:	The recombinant mouse Dpp4 consists of 992 amino acids with a predicted molecular mass of 112.9 kDa.		
Purity:	> 95 % as determined by SDS-PAGE.		
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method		
Biological Activity:	Testing in progress		
Predicted N-terminal:	Glu		
Reconstitution:	Centrifuge vial prior to opening. Add sterile distilled water to a concentration of 0.1 mg/mL and gently pipette the solution up and down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution.		
Storage & Stability:	Stable for up to 1 year from date of receipt at -20°C to -80°C After reconstitution, store working aliquots at -20°C to -80°C. Avoid repeated freeze-thaw cycles.		



SDS-PAGE



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246
Phone: 978-572-1070
Fax: 978-992-0298

E-mail: info@cellsciences.com
Website: www.cellsciences.com