

Igfbp7

Recombinant Mouse IGF-Binding Protein 7 (Fc Tag)

Catalog No.	CRM742A-Fc CRM742B-Fc	Quantity:	20 µg 100 µg
Alternate Names:	Insulin-like growth factor binding protein 7		
Description:	Insulin like growth factor binding protein 7 (IGFBP7) is a member of the IGFBP family. It has been identified in colorectal adenocarcinoma (CRC) cell lines. The Insulin-like growth factor-binding protein also known as IGFBP serves as a carrier protein for Insulin-like growth factor 1. IGFBPs are clearly distinct but are sharing regions with strong homology. All members of the IGFBP family bind IGF-I and IGF-II with about equal affinity. Insulin-like growth factor (IGF) binding proteins (IGFBPs) have been shown to either inhibit or enhance the action of IGF, or act in an IGF-independent manner in the prostate. IGFBP7 could inhibit cell growth, decrease soft agar colony formation activity and induce apoptosis in RKO and SW62 cells. There is mounting evidence that the structure of the IGFBP proteins plays a key role in the regulation of IGF bioavailability, by modulating its molecular size, capillary membrane permeability, target tissue specificity, cell membrane adherence and IGF affinity.		
UniProt ID:	A0A5E1		
Accession Number:	AAH92538.1		
Protein Construction:	A DNA sequence encoding the mouse Igfbp7 (Met1-Leu281) was expressed with the Fc region of human IgG1 at the C-terminus.		
Source:	HEK293 Cells		
Formulation:	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.		
Molecular Weight:	The recombinant mouse Igfbp7 consists of 497 amino acids with a predicted molecular mass of 53.3 kDa.		
Purity:	> 95 % as determined by SDS-PAGE.		
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method		
Biological Activity:	Testing in progress		
Predicted N-terminal:	Ser 26		
Reconstitution:	Centrifuge vial prior to opening. Add sterile distilled water to a concentration of 0.1 mg/mL and gently pipette the solution up and down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution.		
Storage & Stability:	Stable for up to 1 year from date of receipt at -20°C to -80°C After reconstitution, store working aliquots at -20°C to -80°C. Avoid repeated freeze-thaw cycles.		



Cell Sciences®

65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246


Phone: 978-572-1070

Fax: 978-992-0298

E-mail: info@cellsciences.com

Website: www.cellsciences.com

SDS-PAGE



NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246
Phone: 978-572-1070
Fax: 978-992-0298

E-mail: info@cellsciences.com
Website: www.cellsciences.com