

## Tnf



# Recombinant Mouse Tumor Necrosis Factor alpha

<b>Catalog No.</b>	CRT192A CRT192B CRT192C CRT192D	<b>Quantity:</b>	5 µg 20 µg 1.0 mg 100 µg
<b>Alternate Names:</b>	TNF-alpha, Tumor necrosis factor ligand superfamily member 2, TNFSF2, Cachectin		
<b>Description:</b>	Tumor necrosis factor alpha (TNF-alpha) is a multifunctional proinflammatory cytokine secreted by macrophages, monocytes, neutrophils, T cells, NK-cells following their stimulation by bacterial LPS. TNF-alpha activates signals through two receptors, TNFR1, which is expressed on most cell types, and TNFR2, which is expressed mainly on immune cells. TNF-alpha can have many functions, including stimulation of phagocytosis in macrophages, chemoattraction of neutrophils, increase in insulin resistance, and inducing fever, cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation.		
<b>Gene ID:</b>	21926		
<b>UniProt ID:</b>	P06804		
<b>Source:</b>	<i>E. coli</i>		
<b>Molecular Weight:</b>	Monomer, 17.4 kDa (157 aa)		
<b>Formulation:</b>	Lyophilized from a sterile filtered solution containing 10 mM sodium phosphate, 50 mM sodium chloride, pH 7.5.		
<b>Purity:</b>	≥95% by reducing and non-reducing SDS-PAGE		
<b>Endotoxin Level:</b>	≤ 1 EU/µg by kinetic LAL analysis		
<b>Biological Activity:</b>	ED <sub>50</sub> ≤ 20 pg/ml, determined by cytolysis of mouse L929 cells in the presence of Actinomycin D.		
<b>Specific Activity:</b>	≥ 1.0 × 10 <sup>7</sup> U/mg		
<b>Amino Acid Sequence:</b>	MLRSSSQNSS DKPVAHVVAN HQVEEQLEWL SQRANALLAN GMDLKDNQLV VPADGLYLVY SQVLFGQGQC PDYVLLTHTV SRFAISYQEK VNLLSAVKSP CPKDTPEGAE LKPWYEPIYL GGVFQLEKGD QLSAEVNLPK YLDFAESGQV YFGVIAL		
<b>Reconstitution:</b>	<b>Centrifuge vial prior to opening.</b> Add sterile distilled water to a concentration of 0.1 mg/mL and gently pipette the solution up and down the sides of the vial. <b>DO NOT VORTEX.</b> Allow several minutes for complete reconstitution.		



## Storage & Stability:

Store as supplied at -20 °C to -80 °C for up to one year. Upon reconstitution the preparation is stable for up to one month at 2- 8 °C. For long term, reconstitute in working aliquots containing 0.1 % BSA and store at -80 °C. **Avoid repeated freeze-thaw cycles.**



### Mouse TNF-a Gel

Figure: 1 ug in each lane (-) non-reducing conditions and (+) reducing conditions in a 4-20% Tris-Glycine gel, stained with Coomassie Blue. Mouse TNF alpha has a predicted MW of 17.4 kDa.

**NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.**



**Cell Sciences®**  
65 Parker Street  
Unit 11  
Newburyport, MA 01950

Toll Free: 888-769-1246  
Phone: 978-572-1070  
Fax: 978-992-0298

E-mail: [info@cellsciences.com](mailto:info@cellsciences.com)  
Website: [www.cellsciences.com](http://www.cellsciences.com)