## cellsciences.com

## **AREG**

## Recombinant Human Amphiregulin, Animal Free

Catalog No. CS273A-AF Quantity: 10 μg

CS273B-AF 50 μg CS273C-AF 1 mg

Alternate Names: AREG, Colorectum cell-derived growth factor, CRDGF, Schwannoma-derived growth

factor, SDGF

**Description:** Amphiregulin is an EGF related growth factor that signals through the EGF/TGF-a

receptor, and stimulates growth of keratinocytes, epithelial cells and some fibroblasts.

Amphiregulin also inhibits the growth of certain carcinoma cell lines.

Synthesized as a transmembrane protein, Amphiregulin's extracellular domain is

proteolytically processed to release the mature protein. There are 6 conserved cysteine residues, which form 3 intramolecular disulfide bonds essential for biological activity.

Gene ID: 374

UniProt ID: P15514

Source: E. coli

Manufactured without Animal-derived products, in an Animal Free facility.

Molecular Weight: 10.1 kDa, (87 aa) monomer

**Formulation:** Lyophilized from a 0.2 µm filtered solution containing 10 mM sodium phosphate, pH 7.5.

**Purity:** >95% by reducing and non-reducing SDS-PAGE

**Endotoxin Level:**  $\leq 1 \text{ EU/}\mu\text{g}$  determined by kinetic LAL analysis.

**Biological Activity:**  $ED_{50} \le 20$  ng/ml, determined by proliferation of murine 3T3 cells.

**Specific Activity:**  $\geq 5.0 \times 10^4$  Units/mg.

Amino Acid Sequence: SVRVEQVVKP PQNKTESENT SDKPKRKKKG GKNGKNRRNR KKKNPCNAEF

QNFCIHGECK YIEHLEAVTC KCQQEYFGER CGEKSMK

**Reconstitution:** Centrifuged prior to opening. Reconstitute in sterile distilled water or aqueous buffer to

a concentration of 0.1-1.0 mg/mL.

**Storage & Stability:** Store as supplied at -20°C to -80°C for up to 1 year. Upon reconstitution, prepare

working aliquots and store at -20°C to -80°C. It is recommended that a carrier protein

E-mail: info@cellsciences.com

Website: <u>www.cellsciences.com</u>

such as 0.1% HSA or BSA is added for long term storage.


Toll Free: 888-769-1246


Phone: 978-572-1070

Fax: 978-992-0298

Avoid repeated freeze-thaw cycles.

## cellsciences.com





Human Amphiregulin Gel
Figure: 1 ug in each lane (-) non-reducing conditions and (+) reducing conditions in a 4-20% Tris-Glycine gel, stained with Coomassie Blue. Human AREG has a predicted MW of 10.1 kDa.

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

Toll Free: 888-769-1246 Phone: 978-572-1070 Fax: 978-992-0298 E-mail: <u>info@cellsciences.com</u>
Website: <u>www.cellsciences.com</u>