


IL17f

Recombinant Mouse Interleukin-17F, Animal Free

Catalog No.	CS321A-AF CS321B-AF CS321C-AF CS321D-AF	Quantity:	5 µg 25 µg 1 mg 100 µg
Alternate Names:	IL-17F		
Description:	Interleukin 17F (IL-17F) is one of six members of the IL-17 family (IL-17A-F) secreted by activated CD4+ T cells and monocytes. Similar to IL-17A, IL-17F binds to the IL17 RC receptor and promotes the production of IL-6, IL-8, GM-CSF and increases matrix turnover rates. IL-17F is also thought to inhibit angiogenesis and induce endothelial cells to produce IL-2, MCP-1 and TGF-β1.		
Gene ID:	257630		
UniProt ID:	Q7TNI7		
Source:	<i>E. coli</i> Manufactured without Animal-derived products, in an Animal Free facility.		
Molecular Weight:	15.0/30.0 kDa (134/268 aa) dimer		
Formulation:	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)		
Purity:	≥ 95% determined by reducing and non-reducing SDS-PAGE		
Endotoxin Level:	≤ 1 EU/µg by kinetic LAL analysis		
Amino Acid Sequence:	MRKNPKAGVP ALQKAGNCPD LEDNTVRVDI RIFNQNQGSI VPREFQNRSS SPWDYNITRD PHRFPSIEAI AQCRHSGCIN AQQQEDSTMN SVAIQQEILV LRREPQGCSN SFRLEKMLLK VGCTCVKPIV HQAA		
Reconstitution:	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at a concentration of 0.1 mg/mL, which can be further diluted into other aqueous solutions.		
Storage & Stability:	Lyophilized product is stable at room temperature for shipping purposes. Upon receipt, store at -20°C to -80°C for up to 1 year. Upon reconstitution, the preparation is stable for up to one month at 2-8°C. For long term storage, prepare working aliquots and store at -20 to -80°C. For maximal stability, dilute to working aliquots in a 0.1% BSA solution. Avoid repeated freeze-thaw cycles.		



Mouse IL-17F Gel

Figure: 1 ug run under (-) non-reducing conditions and (+) reducing conditions in a 4-20% Tris-Glycine gel, stained with Coomassie Blue. Mouse IL-17F is a homodimer with a predicted MW of 30.0 kDa (each monomer is 15.0 kDa).

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences®
65 Parker Street
Unit 11
Newburyport, MA 01950

Toll Free: 888-769-1246
Phone: 978-572-1070
Fax: 978-992-0298

E-mail: info@cellsciences.com
Website: www.cellsciences.com