

BLK Recombinant Human B Lymphoid Tyrosine Kinase (Met₁-Pro₅₀₅) Active GST-His

Catalog No. CSI11173 Quantity: 50 µg

Alternate Names: MGC10442, BLK nonreceptor tyrosine kinase

Description: Human BLK, Amino acids M₁-P₅₀₅ (as in GenBank entry NM_001715.2)*, N-terminally

fused to GST-HIS₆-Thrombin cleavage site

*Sequence may contain documented polymorphisms

Detailed sequence on request

Concentration: 0.120 μg/μl

Gene ID: 640

Protein Accession No: NM_001715.2

Source: Baculovirus infected Sf9 cells

Molecular Weight: Theoretical MW_{Eusion Protein}: 87,102 Da

Formulation: 50 mM Tris-HCl, pH 8.0 + 100 mM NaCl + 5 mM DTT + 4 mM reduced glutathione, 20%

glycerol

Purification: One-step affinity purification using GSH-agarose

Product Identity: BLK was confirmed as human BLK by mass spectroscopy LC-ESI-MS/MS

Specific Activity: 167 pmol/µg×min

Method for determination of K_m value and specific activity:

· Assay conditions:

60 mM HEPES-NaOH, pH 7.5

3 mM MgCl₂ 3 mM MnCl₂

3 µM Na-orthovanadate

1.2 mM DTT

2.5 μg / 50 μl PEG_{20.000}

ATP (variable)


Substrate: Poly(Glu,Tyr) $_{4:1}$ (Sigma P-0275), 1,0 μ g / 50 μ l Recombinant BLK: 200 μ g / 50 μ l

Filter binding assay

MSFC membrane (Millipore)

Storage & Stability: Store in working aliquots at -80°C. Avoid repeated freeze-thaw cycles.

Fax: 781-828-0542



Toll Free: 888-769-1246 E-mail: info@cellsciences.com
Phone: 781-828-0610 Website: www.cellsciences.com

cellsciences.com

Determination of K_m value for ATP: Western blot analysis: Coomassie stain: 200 anti-GST kDa V (pmol ATP/µg×min) 150-150 -250 150 100 100 75-GST-BLK V_{Max} = 167 pmol/µg×min 50 -37 31-50 100 150 200 0 Conc. ATP (µM) 2.0 µg GST-BLK 600 ng OST-BLK

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

Toll Free: 888-769-1246 E-mail: info@cellsciences.com
Phone: 781-828-0610 Website: www.cellsciences.com
Fax: 781-828-0542