

Recombinant HSV/Herpes Simplex Virus-2 gD (aa 266-394) GST

Catalog No. CSI15834A Quantity: 100 µg

CSI15834B 0.5 mg CSI15834C 1.0 mg

Description: Entry of HSV into the host cell involves interactions of several viral glycoproteins with cell

surface receptors. The virus particle is covered by an envelope which, when bound to specific receptors on the cell surface, will fuse with the cell membrane and create an opening, or pore, through which the virus enters the host cell. The sequential stages of HSV entry are analogous to those of other viruses. At first, complementary receptors on the virus and cell surface bring the two membranes into proximity. In an intermediate state, the two membranes begin to merge, forming a hemifusion state. Finally, a stable entry pore is formed through which the viral envelope contents are introduced to the host

cell.

The *E. coli* derived 39.7 kDa recombinant protein contains the HSV-2 gD immunodominant regions 266-394 amino acids, fused with 26 kDa GST-tag.

Source: E. coli

Formulation: 25 mM Tris-HCl + 1 mM EDTA + 50% glycerol.

Purity: HSV-2 gD protein is >95% pure as determined by 10% PAGE (coomassie staining).

Purification Method: HSV-2 gD protein was purified by proprietary chromatographic technique.

Specific Activity: Immunoreactive with sera of HSV-infected individuals.

Storage & Stability: HSV-2 gD Protein although stable at 4°C for 1 week, should be stored below -18°C.

Toll Free: 888-769-1246

Phone: 781-828-0610

Fax: 781-828-0542

E-mail: <u>info@cellsciences.com</u>
Website: www.cellsciences.com

Please prevent freeze thaw cycles.

Applications: HSV-2 gD antigen is suitable for ELISA and Western blots, excellent antigen for detection

of HSV with minimal specificity problems.

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.