

Recombinant Herpes Simplex Virus-8 Mosaic GST

Catalog No. CSI15837A Quantity: 100 µg

CSI15837B 0.5 mg CSI15837C 1.0 mg

Description: Entry of HSV into the host cell involves interactions of several viral glycoproteins with cell

surface receptors. The virus particle is covered by an envelope which, when bound to specific receptors on the cell surface, will fuse with the cell membrane and create an opening, or pore, through which the virus enters the host cell. The sequential stages of HSV entry are analogous to those of other viruses. At first, complementary receptors on the virus and cell surface bring the two membranes into proximity. In an intermediate state, the two membranes begin to merge, forming a hemifusion state. Finally, a stable entry pore is formed through which the viral envelope contents are introduced to the host

cell.

The *E. coli* derived recombinant protein contains the C-terminal immunodominant regions from ORF65 140-170 a.a. and N-terminal regions from ORF8 32-62 a.a. The protein is

fused with a GST tag.

Source: E. coli

Formulation: 100 mM NaCl + 0.1% SDS and 50% glycerol.

Purity: HSV-8 Mosaic protein is >95% pure as determined by 10% PAGE (coomassie staining).

Purification Method: HSV-8 Mosaic was purified by proprietary chromatographic technique.

Specific Activity: Immunoreactive with sera of HSV-8 infected individuals.

Storage & Stability: HSV-8 Mosaic protein although stable at 4°C for 1 week, should be stored below -18°C.

Please prevent freeze thaw cycles.

Applications: HSV-8 Mosaic antigen is suitable for ELISA and Western blots, excellent antigen for

Toll Free: 888-769-1246

Phone: 781-828-0610

Fax: 781-828-0542

E-mail: info@cellsciences.com

Website: www.cellsciences.com

detection of HSV with minimal specificity problems.

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.