

TACI Peptide

Catalog No:	PX242BP	Size:	50 µg
Sequence:	NH ₂ -SGLGRSRRGGRSRVD-COOH		
Location:	Amino acids 2 TO 16 of human TACI (1).		
Application:	The peptide is used for blocking the antibody activity of TACI (catalog number: PX242). It usually blocks the antibody activity completely in Western blot by incubating the peptide with equal volume of antibody for 30 min at 37°C.		
Storage:	It is supplied as 50 µg at 200 µg/ml in PBS pH7.2 (10 mM NaH ₂ PO ₄ , 10 mM Na ₂ HPO ₄ , 130 mM NaCl) containing 0.1% bovine serum albumin and 0.02% sodium azide. Stable for one year if stored at -20°C.		
Reference:	1. Von Bulow GU, Bram RJ. NF-AT activation induced by a CAML-interacting member of the tumor necrosis factor receptor superfamily. <i>Science</i> . 1997; 278 (5335):138-41.		

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Cell Sciences, Inc.
480 Neponset Street
Building 12A
Canton, MA 02021

Toll Free: 888 769-1246
Phone: 781 828-0610
Fax: 781 828-0542

E-mail: info@cellsciences.com
Web Site: www.cellsciences.com