

## **Rat Small Intestine Tissue Lysate**

Catalog No. PX419A Quantity: 100 µg

**Description:** Rat small Intestine tissue lysate was prepared by homogenization in modified RIPA

buffer (150 mM sodium chloride, 50 mM Tris-HCl, pH 7.4, 1 mM

ethylenediaminetetraacetic acid, 1 mM phenylmethylsulfonyl fluoride, 1% Triton X-100, 1% sodium deoxycholic acid, 0.1% sodium dodecylsulfate, 5  $\mu$ g/ml of aprotinin, 5  $\mu$ g/ml of leupeptin. Tissue and cell debris was removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay. The product was boiled for 5 min in 1 x SDS

sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5%  $\beta$ -mercaptoethanol.

**Source:** Rat small intestine tissue from white laboratory rats

Concentration: 2 mg/ml

Formulation: Small Intestine tissue lysate is supplied in SDS sample buffer containing 5% β-

mercaptoethanol.

Species: Rat

Applications: Centrifuge vial before use. Rat small Intestine tissue lysate is ready to load on SDS-

PAGE for Western blotting, 10-20 µg per lane is recommended for mini gel.

**Storage & Stability:** Store at 2-8°C for continuous use. Store in working aliquots at -80°C for up to 1 year.

Toll Free: 888-769-1246

Phone: 978-572-1070

Fax: 978-992-0298

E-mail: info@cellsciences.com

Website: www.cellsciences.com

Avoid repeated freeze-thaw cycles.

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.