

Rat Testis Tissue Lysate

Catalog No. PX422A **Quantity:** 100 µg

Description: Rat testis tissue lysate was prepared by homogenization in modified RIPA buffer (150 mM sodium chloride, 50 mM Tris-HCl, pH 7.4, 1 mM ethylenediaminetetraacetic acid, 1 mM phenylmethylsulfonyl fluoride, 1% Triton X-100, 1% sodium deoxycholic acid, 0.1% sodium dodecylsulfate, 5 µg/ml of aprotinin, 5 µg/ml of leupeptin. Tissue and cell debris was removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay. The product was boiled for 5 min in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% β-mercaptoethanol.

Source: Rat testis tissue from outbred Sprague Dawley Albino rat

Concentration: 2 mg/ml

Formulation: Testis tissue lysate is supplied in SDS sample buffer containing 5% β-mercaptoethanol.

Species: Rat

Applications: **Centrifuge vial before use.** Rat testis tissue lysate is ready to load on SDS-PAGE for Western blotting. It is recommended to load 10 µg to 20 µg per lane for mini gel.

Storage & Stability: Store at 2-8°C for continuous use. Store in working aliquots at -80°C for up to 1 year. **Avoid repeated freeze-thaw cycles.**

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.