


Data Sheet

Product Name:	Ethyl gallate
Cat. No.:	CS-0009060
CAS No.:	831-61-8
Molecular Formula:	C ₉ H ₁₀ O ₅
Molecular Weight:	198.17
Target:	Bacterial
Pathway:	Anti-infection
Solubility:	DMSO : 150 mg/mL (756.93 mM; Need ultrasonic)



BIOLOGICAL ACTIVITY:

Ethyl gallate is a nonflavonoid phenolic compound and also a scavenger of hydrogen peroxide. **In Vitro:** Ethyl gallate is a nonflavonoid phenolic compound and also a scavenger of hydrogen peroxide. After treatment for 24 h or 48 h with Ethyl gallate, HL-60 cells show changes in morphology, including shrinkage of the cell membrane and the development of apoptotic bodies. Consistent with these effects, the viability of Ethyl gallate-treated cells decreases in a time- and dose-dependent manner, demonstrating that Ethyl gallate has a cytotoxic effect on HL-60 cells. Ethyl gallate treatment increases the proportion of cells in subG1 phase in a concentration- and time-dependent manner. Treatment of cells for 24 h or 48 h with 50 μ M or 75 μ M Ethyl gallate increases the percentage of cells in the subG1 phase from a baseline of 2.9% to 26.5% or 52.6%, respectively. It is found that Ethyl gallate treatment of HL-60 cells decreases the expression of Bcl-2 at 75 μ M Ethyl gallate, and increases Bax and truncated Bid (tBid) expression at 24 h^[1]. **In Vivo:** No significant difference in the serum total protein, albumin, globulin and glucose is found between the rats fed with *A. nilotica* (L.) leaf extract on ethyl gallate equivalent basis and those fed with Ethyl gallate alone. Significant differences in total bilirubin level, however, exist between the rats that receive *A. nilotica* (L.) leaf extract, 500 mg/kg body weight (ethyl gallate equivalent of 10 mg/kg, 0.34 \pm 0.01 mg/dL) and those receiving 10 mg/kg body weight of Ethyl gallate (0.26 \pm 0.01 mg/dL). Significant difference is found for ALT between groups fed with 500 and 1000 mg/kg body weight of *A. nilotica* (L.) leaf extract (26.52 \pm 1.23 and 30.05 \pm 1.38 U/L) and 10 and 20 mg/kg of Ethyl gallate (20.50 \pm 0.94 and 24.67 \pm 1.13 U/L)^[2].

PROTOCOL (Extracted from published papers and Only for reference)

Kinase Assay: ^[1]The expression of apoptosis-related proteins (caspases-8, -9, -3; AIF; Endo G; Bid; Bax; and Bcl-2) in HL-60 cells is determined by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) of lysates followed by western blotting. For this, HL-60 cells (1.5 \times 10⁶) are treated with **50 μ M or 75 μ M Ethyl gallate for 6 h, 12 h, or 24 h**. Total cell lysates are obtained by resuspending cells in ice-cold radioimmunoprecipitation assay (RIPA) buffer for 30 min followed by centrifugation. Protein concentration is determined using a NanoDrop spectrophotometer. Aliquots of lysates (100 μ g protein equivalents) are resolved by 12% SDS-PAGE and transferred onto nitrocellulose membranes^[1]. **Cell Assay:** ^[1]**HL-60 cells (1 \times 10⁶) are treated with 50 μ M or 75 μ M Ethyl gallate for 24 h or 48 h at 37°C.** Cells are then harvested by centrifugation and fixed in 70% ethanol at 4°C for 24 h. Fixed cells are resuspended in PBS containing 40 μ g/mL Propidium iodide (PI), 100 μ g/mL RNase A, and 0.1% Triton X-100 and incubated in the dark for 30 min at room temperature. Cell cycle distribution is analyzed by flow cytometry on a FACSCalibur. To investigate apoptotic cells, HL-60 cells (1 \times 10⁶) incubated with different concentration of **50 μ M, 75 μ M and 100 μ M Ethyl gallate for 24 h or 48 h at 37°C**, and then DAPI staining is conducted. The cells are photographed using a fluorescence microscopy^[1]. **Animal Administration:** ^[2]Forty eight **female albino Wistar rats** of six to eight weeks old are used and divided into eight groups based on their body weights. Group 1 rats serve as control receiving 1.0 mL of the vehicle (0.1% ethanol); Group 2 rats receive *A. nilotica* (L.) leaf extract (250 mg/kg body weight); Group 3 rats receive *A. nilotica* (L.) leaf extract (500 mg/kg body weight); Group 4 rats receive *A. nilotica* (L.) leaf extract (1000 mg/kg body weight); Group 5 rats receive *A. nilotica* (L.) leaf extract (2000 mg/kg body weight); Group 6 rats receive **Ethyl gallate (5**

mg/kg body weight); Group 7 rats receive **Ethyl gallate (10 mg/kg body weight)**; Group 8 rats receive **Ethyl gallate (20 mg/kg body weight)**. Body weights are recorded on 0th and 14th day for each group and all rats are decapitated after an overnight fast^[2].

References:

[1]. Kim WH, et al. Ethyl gallate induces apoptosis of HL-60 cells by promoting the expression of caspases-8, -9, -3, apoptosis-inducing factor and endonuclease G. *Int J Mol Sci.* 2012;13(9):11912-22.

[2]. Mohan S, et al. In vitro protection of biological macromolecules against oxidative stress and in vivo toxicity evaluation of *Acacia nilotica* (L.) and ethyl gallate in rats. *BMC Complement Altern Med.* 2014 Jul 21;14:257.

CAIndexNames:

Benzoic acid, 3,4,5-trihydroxy-, ethyl ester

SMILES:

O=C(OCC)C1=CC(O)=C(O)C(O)=C1

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 732-484-9848 Fax: 888-484-5008 E-mail: sales@ChemScene.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA