


Data Sheet

Product Name:	Syringaldehyde
Cat. No.:	CS-0016810
CAS No.:	134-96-3
Molecular Formula:	C ₉ H ₁₀ O ₄
Molecular Weight:	182.17
Target:	COX
Pathway:	Immunology/Inflammation
Solubility:	10 mM in DMSO



BIOLOGICAL ACTIVITY:

Syringaldehyde is a polyphenolic compound belonging to the group of flavonoids and is found in different plant species like *Manihot esculenta* and *Magnolia officinalis*^[1]. Syringaldehyde moderately inhibits **COX-2** activity with an **IC₅₀** of 3.5 µg/mL^[2]. Anti-hyperglycemic and anti-inflammatory activities^[1]. **In Vitro:** Syringaldehyde inhibits COX-2 activity in a dose-dependent manner with an **IC₅₀** of 3.5 µg/mL^[2]. **In Vivo:** Syringaldehyde exerts anti-hyperglycemic effect in rat model of diabetes induced by streptozotocin. Apart from antioxidant capability, Syringaldehyde also has anti-inflammatory activity as it is found to have inhibitory action on cyclooxygenase 2 (COX-2) in mouse macrophage cell line^[1].

PROTOCOL (Extracted from published papers and Only for reference)

Animal Administration: Rats^[1]

Adult male albino wistar rats (weighing 180-220 g)

12.5 mg/kg, 25 mg/kg, and 50 mg/kg; p.o.; for 21 days

Adult male albino wistar rats are randomly divided into six groups (n=6), first group serves as control and receives vehicle (orally) for 21 days. Second group is given Syringaldehyde in saline orally at 50 mg/kg for 21 days. Third group receives vehicle for a period of 21 days and then ISO (100 mg/kg, s.c.) on 20th and 21st day at an interval of 24 h. Fourth group is given Syringaldehyde for 21 days at 12.5 mg/kg, p.o. and ISO on 20th and 21st day. Fifth group receives Syringaldehyde at concentration of 25 mg/kg, p.o. for 21 days and ISO on 20th and 21st day. The sixth group is treated with 50 mg/kg of Syringaldehyde for 21 days and ISO on 20th and 21st day. During the experimental procedure body weight of animals are monitored and on 22nd day, 24 h after second injection of ISO, rats are sacrificed by cervical decapitation.

Note: Body weight of animals belonging to different groups did not differ significantly but heart weight of ISO challenged animals were highly elevated ($p \leq 0.05$) than control rats. However, treatment of rats with Syringaldehyde significantly reduced heart weight in a dose-dependent way. The rats treated with Syringaldehyde alone displayed an insignificant difference from control group.

References:

[1]. Shahzad S, et al. Protective effect of syringaldehyde on biomolecular oxidation, inflammation and histopathological alterations in isoproterenol induced cardiotoxicity in rats. *Biomed Pharmacother.* 2018 Dec;108:625-633.

[2]. Stanikunaite R, et al. Cyclooxygenase-2 inhibitory and antioxidant compounds from the truffle *Elaphomyces granulatus*. *Phytother Res.* 2009 Apr;23(4):575-8.

CAIndexNames:

Benzaldehyde, 4-hydroxy-3,5-dimethoxy-

SMILES:

O=CC1=CC(OC)=C(O)C(OC)=C1

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 732-484-9848 Fax: 888-484-5008 E-mail: sales@ChemScene.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA