

Data Sheet

Product Name:	Adrenocorticotrophic Hormone (ACTH) (1-39), human(TFA)
Cat. No.:	CS-0062643
Molecular Formula:	C207H308N56O58S.C2HF3O2
Molecular Weight:	4655.16
Target:	Others
Pathway:	Others
Solubility:	H2O

SYSMEHFRWG KPVGKKRRPV KVYPNGAEDE SAEAPPLEF



BIOLOGICAL ACTIVITY:

Adrenocorticotrophic Hormone (ACTH) (1-39), human(TFA) is a **melanocortin receptor** agonist. IC50 & Target: Melanocortin receptor^[1]
In Vitro: Adrenocorticotrophic Hormone (ACTH) (1-39), human (ACTH 1-39), a member of the melanocortin family, stimulates production of CS by the adrenals, but melanocortin receptors are also found in the central nervous system (CNS) and on immune cells. ACTH 1-39 protects neurons in vitro from several apoptotic, excitotoxic and inflammation-related insults^[1]. The conditioned medium (CM) is prepared from untreated astroglia (AS) cultures and from AS cultures treated with 200 nM ACTH 1-39 for 24 h, washed to remove ACTH 1-39, then incubated for another 24 h in DMEM. In initial experiments, no difference is found in oligodendroglia (OL) viability in the presence of OL defined medium with 2% newborn calf serum (NCS) or AS CM (prepared in DMEM with no serum). After 24 h, OL death under each condition varies between 1 and 4%. Similar results for OL viability are obtained with microglia (MG) CM. In subsequent experiments, controls in each experiment consist of OL in defined medium with 2% NCS^[2].

PROTOCOL (Extracted from published papers and Only for reference)

Cell Assay: ^[2]For analysis of the effects of ACTH 1-39 or CM on **oligodendroglia (OL)** death, purified OL cultures are incubated with **ACTH 1-39** at **200 nM** or the various CM for 30 min before addition of the toxic agents. Cell death is assessed after 1 day using trypan blue uptake as the indicator of cell death. Trypan blue is considered a preferred method for measurement of total cell death compared to terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL), which measures only apoptosis, or live/dead fluorescent assays, which may not detect permeable dead cells with degraded DNA, thus underestimating cell death. Differentiated OL are identified by their characteristic morphology, that is, rounded or oval birefringent cells with multiple lacy branching processes, and in some cases by immunostaining with antibodies to galactolipids^[2].

References:

- [1]. Lisak RP, et al. Melanocortin receptor agonist ACTH 1-39 protects rat forebrain neurons from apoptotic, excitotoxic and inflammation-related damage. *Exp Neurol*. 2015 Nov;273:161-7.
- [2]. Lisak RP, et al. The melanocortin ACTH 1-39 promotes protection of oligodendrocytes by astroglia. *J Neurol Sci*. 2016 Mar 15;362:21-6.

CAIndexNames:

Adrenocorticotrophic Hormone (ACTH) (1-39), human(TFA)

SMILES:

OC(C(F)(F)F)=O.[SYSMEHFRWG KPVGKKRRPV KVYPNGAEDE SAEAPLEF]

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 732-484-9848 Fax: 888-484-5008 E-mail: sales@ChemScene.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA