


Data Sheet

| | |
|---------------------------|---|
| Product Name: | Tenovin-6 (Hydrochloride) |
| Cat. No.: | CS-2321 |
| CAS No.: | 1011301-29-3 |
| Molecular Formula: | C ₂₅ H ₃₅ ClN ₄ O ₂ S |
| Molecular Weight: | 491.09 |
| Target: | Autophagy; MDM-2/p53; Sirtuin |
| Pathway: | Apoptosis; Autophagy; Cell Cycle/DNA Damage; Epigenetics |
| Solubility: | DMSO : ≥ 49 mg/mL (99.78 mM) |



BIOLOGICAL ACTIVITY:

Tenovin-6 Hydrochloride is an inhibitor of **SIRT1** and **SIRT2**, slightly inhibits **HDAC8**, and is also a potent activator of **p53**, with **IC₅₀s** of 21 μ M, 10 μ M, and 67 μ M for SirT1, SirT2, and SirT3, respectively. **IC₅₀ & Target:** IC₅₀: 21 μ M (SirT1), 10 μ M (SirT2), 67 μ M (SirT3)^[1] **In Vitro:** Tenovin-6 inhibits the growth of *S. cerevisiae* cultures with an IC₅₀ of 30 μ M and is more toxic to yeast than the less water-soluble tenovin-1. Tenovin-6 rapidly increases the levels of endogenous K382-Ac p53 in MCF-7 cells^[1]. Tenovin-6 (0 to 15 μ M) dose dependently increases the level of LC3-II in diverse cell types, and the increase is ATG5/7 dependent. Tenovin-6 treatment also increases the number and intensity of autophagic vesicles with or without the presence of Torin 1, and prevents Torin 1-induced SQSTM1/p62 degradation. Tenovin-6 affects the acidification of autolysosomes and impairs the hydrolytic activity of lysosomes but does not affect the fusion between autophagosomes and lysosomes. That tenovin-6 inhibits autophagy does not correlate with p53 activation and SIRT1/2 inhibition by knockdown or knockout cannot mimic the effect of tenovin-6 on LC3B accumulation^[2]. Tenovin-6 (0, 1, 2.5, 5 or 10 μ M) potently inhibits cell proliferation in a dose- and time-dependent manner in all OCI-Ly1, DHL-10, U2932, RIVA, HBL1 and OCI-Ly10 cell lines. Tenovin-6 consistently increases LC3B-II level in DLBCL cell lines by inhibiting the classical autophagy pathway, without activating p53, and the increase is independent of SIRT1/2/3 and p53. Tenovin-6 induces apoptosis through the extrinsic cell-death pathway^[3]. Tenovin-6 suppresses the growth of UM cells with IC₅₀ of 12.8 μ M, 11.0 μ M, 14.58 μ M and 9.62 μ M for 92.1, Mel 270, Omm 1 and Omm 2.3 cells, respectively^[4]. **In Vivo:** Tenovin-6 (50 mg/kg, i.p.) inhibits the growth of tumor in mice^[1].

PROTOCOL (Extracted from published papers and Only for reference)

Kinase Assay: ^[1]Assays are carried out using purified components in the Fluor de Lys Fluorescent Assay Systems. Relevant FdL substrates are used at 7 μ M and NAD⁺ at 1 mM. Tenovins are solubilized in DMSO with the final DMSO concentration in the reaction being less than 0.25%. For SirT1 and HDAC8, one unit of enzyme is used per reaction, and for SirT2 and SirT3, five units is used per reaction. Reactions are carried out at 37°C for 1 hr. **Cell Assay:** ^[4]The MTS assay is used to evaluate cell viability. UM cells are seeded into each well of 96-well plates (5,000 cells/well) and treated the next day with control or Tenovin-6 in an increasing concentrations from 0 to 20 μ M for 68 h, and then MTS is added at 20 μ L/well to be read at a wave length of 490 nm, the IC₅₀ is determined by curve fitting of the sigmoidal dose-response curve. **Animal Administration:** Tenovin-6 is formulated in vehicle solution containing cyclodextrin 20% (w/v) and DMSO 10% (v/v).^[1]Female SCID mice are injected subcutaneously with 1×10⁶ ARN8 cells suspended in matrigel. Tumors are allowed to reach a size of approximately 10 mm³. Tenovin-6 is administered daily at 50 mg/kg by intraperitoneal injection. Control animals are treated with vehicle solution containing cyclodextrin 20% (w/v) and DMSO 10% (v/v). Tumor diameters are measured using calipers, and volumes are calculated using the equation $V = \pi d_1^2 d_2 / 6$. Median values of tumor size are calculated for each time point as well as the corresponding 95% confidence intervals. Comparison of control and drug-treated tumor size distributions are made by Mann-Whitney U-test. An alpha-level of 0.05 is considered appropriate for determination of statistical significance.

References:

- [1]. Lain S, et al. Discovery, in vivo activity, and mechanism of action of a small-molecule p53 activator. *Cancer Cell*. 2008 May;13(5):454-63.
- [2]. Yuan H, et al. Tenovin-6 impairs autophagy by inhibiting autophagic flux. *Cell Death Dis*. 2017 Feb 9;8(2):e2608.
- [3]. Yuan H, et al. Tenovin-6 inhibits proliferation and survival of diffuse large B-cell lymphoma cells by blocking autophagy. *Oncotarget*. 2017 Feb 28;8(9):14912-14924.
- [4]. Dai W, et al. Class III-specific HDAC inhibitor Tenovin-6 induces apoptosis, suppresses migration and eliminates cancer stem cells in uveal melanoma. *Sci Rep*. 2016 Mar 4;6:22622.

CAIndexNames:

Benzamide, N-[[[4-[[5-(dimethylamino)-1-oxopentyl]amino]phenyl]amino]thioxomethyl]-4-(1,1-dimethylethyl)-, hydrochloride (1:1)

SMILES:

O=C(NC(NC1=CC=C(NC(CCCCN(C)C)=O)C=C1)=S)C2=CC=C(C(C)C)C=C2.Cl

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 732-484-9848 Fax: 888-484-5008 E-mail: sales@ChemScene.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA