


Data Sheet

Product Name:	THZ1
Cat. No.:	CS-3167
CAS No.:	1604810-83-4
Molecular Formula:	C ₃₁ H ₂₈ ClN ₇ O ₂
Molecular Weight:	566.05
Target:	CDK
Pathway:	Cell Cycle/DNA Damage
Solubility:	DMSO : 100 mg/mL (176.66 mM; Need ultrasonic)



BIOLOGICAL ACTIVITY:

THZ1 is a selective and potent covalent **CDK7** inhibitor with an **IC₅₀** of 3.2 nM. THZ1 also inhibits the closely related kinases **CDK12** and **CDK13** and downregulates MYC expression^{[1][2]}. **IC₅₀ & Target:** IC₅₀: 3.2 nM (CDK7)^[1] **In Vitro:** THZ1 inhibits Jurkat cell and Loucy cell with IC₅₀ of 50 nM, and 0.55 nM, respectively. THZ1 (9, 27, 83, 250, 750, and 2500 nM) inhibits CDK12 but at higher concentrations compared to CDK7. THZ1 (1 μM) irreversibly inhibits RNAPII CTD and CAK phosphorylation. THZ1 (2.5 μM) irreversibly inhibits RNAPII CTD phosphorylation by covalently targeting a unique cysteine located outside the kinase domain of CDK7 in Hela S3 cells. THZ1 (250 nM) causes decreased cellular proliferation and an increase in apoptotic index with concomitant reduction in anti-apoptotic proteins, most notably MCL-1 and XIAP in T-ALL cell lines^[1].

All genotypically-distinct human (hSCLC) cell lines exhibit high sensitivity to THZ1, with an IC₅₀ in the range of 5-20 nM^[3]. **In Vivo:** THZ1 (10 mg/kg) demonstrates potent killing of primary chronic lymphocytic leukemia (CLL) cells and anti-proliferative activity against primary TALL cells and in vivo against a human T-ALL xenograft^[1].

THZ1 (10 mg/kg, i.v.) inhibits tumor growth in a mouse model of human MYCN-amplified NB and shows no toxicity^[4].

THZ1 (10 mg/kg, i.p.) completely suppresses oesophageal squamous cell carcinoma tumour growth in vivo without loss of body weight or other common toxic effects^[5].

PROTOCOL (Extracted from published papers and Only for reference)

Kinase Assay: ^[1]For kinase assays following immunoprecipitation of FLAG-CDK7 protein from HCT116 or FLAG-CDK12 from 293A cellular lysates, cells are first treated with THZ1, THZ1-R, or DMSO for 4 hrs at 37°C. Cells are then harvested by lysis in 50 mM Tris HCl pH 8.0, 150 mM NaCl, 1% NP-40, 5 mM EDTA, and protease/phosphatase cocktails. Exogenous CDK7 or CDK12 proteins are immunoprecipitated from cellular lysates using FLAG antibody- conjugated agarose beads. Precipitated proteins are washed with lysis buffer 6 times, followed by 2 washes with kinase buffer (40 mM Hepes pH 7.5, 150 mM NaCl, 10 mM MgCl₂, 5% glycerol) and subjected to in vitro kinase assays at 30°C for 45 minutes using 1 μg of the large subunit of RNAPII (RPB1) as substrate and 25 μM ATP and 10 μCi of 32P ATP. Kinase assays using recombinant CDK7/TFIIH/MAT1 are conducted in the manner as described above using 25 ng of CAK complex per reaction. For kinase assays designed to test time-dependent inactivation of CDK7 kinase activity, CAK complex is pre-incubated with indicated concentrations of THZ1, THZ1-R, or DMSO in kinase buffer without ATP for 4 hrs at 30°C prior to being subjected to kinase assay conditions^[1]. **Cell Assay:** THZ1 is dissolved in DMSO, and then diluted with appropriate media before use^[1].

^[1]Jurkat, Loucy, KOPTK1, and DND-41 cell lines are seeded in 384-well microplates at 15% confluency in medium with 5% FBS and penicillin/streptavidin. Cells are treated with THZ1 (2, 10, 50, 250, 1250, and 6250 nM) or DMSO for 72 hrs and cell viability is determined using resazurin^[1]. **Animal Administration:** THZ1 is prepared in 10 % DMSO in 5 % glucose water^[1].^[1]Mice^[1]

Thirty-two NOD-SCIDIL2Rcynull (NSG) 9-week old female mice are divided into treatment groups based on mean BLI as follows: THZ1 10 mg/kg qD, THZ1 10 mg/kg BID, and vehicle (10% DMSO in D5W) BID (n=10 for all groups). Two mice are excluded, one with the highest and one with the lowest BLI. All treatments are administered via IV injection in the lateral tail vein in a volume of 3.3 μL/g

(non-blinded). Mice are imaged and weighed every 3-5 days. Mice are treated for four weeks and on the final day mice are imaged, dosed and sacrificed approximately 5-6 hrs post dose. Upon sacrifice, blood is collected via cardiac puncture in EDTA tubes; a portion (300 uL) is processed for plasma. Liver and spleen tissues are collected from each mouse with half of each sample flash frozen and half of each sample fixed. Blood plasma and liver samples are processed for pharmacokinetics analysis of THZ1. Spleen tissues are homogenized and lysed and processed for pharmacodynamics analysis of THZ1 target engagement.

References:

- [1]. Kwiatkowski N, et al. Targeting transcription regulation in cancer with a covalent CDK7 inhibitor. *Nature*. 2014 Jul 31;511(7511):616-20.
- [2]. Zeng M, et al. Targeting MYC dependency in ovarian cancer through inhibition of CDK7 and CDK12/13. *Elife*. 2018 Nov 13;7. pii: e39030.
- [3]. Christensen CL, et al. Targeting transcriptional addictions in small cell lung cancer with a covalent CDK7 inhibitor. *Cancer Cell*. 2014 Dec 8;26(6):909-22.
- [4]. Chipumuro, et al. CDK7 inhibition suppresses super-enhancer-linked oncogenic transcription in MYCN-driven cancer. *Cell*. 2014 Nov 20;159(5):1126-39. ?
- [5]. Jiang YY, et al. Targeting super-enhancer-associated oncogenes in oesophageal squamous cell carcinoma. *Gut*. 2016 May 10. pii: gutjnl-2016-311818.
- [6]. Jiang YY, et al. Targeting super-enhancer-associated oncogenes in oesophageal squamous cell carcinoma. *Gut*. 2016 May 10. pii: gutjnl-2016-311818.

CAIndexNames:

Benzamide, N-[3-[[5-chloro-4-(1H-indol-3-yl)-2-pyrimidinyl]amino]phenyl]-4-[[2E)-4-(dimethylamino)-1-oxo-2-buten-1-yl]amino]-

SMILES:

C1C1=CN=C(NC2=CC(NC(C3=CC=C(NC(/C=C/CN(C)C)=O)C=C3)=O)=CC=C2)N=C1C4=CNC5=CC=CC=C54

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 732-484-9848 Fax: 888-484-5008 E-mail: sales@ChemScene.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA