


Data Sheet

Product Name:	Prostaglandin E2
Cat. No.:	CS-6932
CAS No.:	363-24-6
Molecular Formula:	C ₂₀ H ₃₂ O ₅
Molecular Weight:	352.47
Target:	Endogenous Metabolite; Prostaglandin Receptor
Pathway:	GPCR/G Protein; Metabolic Enzyme/Protease
Solubility:	DMSO : ≥ 100 mg/mL (283.71 mM)



BIOLOGICAL ACTIVITY:

Prostaglandin E2 is a vasodilator isolated from prostate gland secretion, working by binding and activating the **prostaglandin E2 receptor**. **In Vitro:** PGE2 shows inhibition of IL 2 production in the mixture of irradiated and nonirradiated T lymphocytes. PGE2 (0.1-10 μM) dose-dependently inhibits the production of IL 2. PGE2 acts during the inductive phase of activation of suppressor cells. Preincubation of T lymphocytes with PGE2 induces cells that suppress IL 2 production and PHA proliferation^[1]. **In Vivo:** PGE2 (0.3 μg/kg, i.p.) significantly reduces the number of peritoneal macrophages undergoing phagocytosis of the methacrylate microbeads in rats^[2]. PGE2 (0.1 mg/min, i.a.) increases renal blood flow. PGE2 produces a biphasic change in renal vascular resistance, vasodilatation starts at 0.01 mg/min and is maximal at about 3 mg/min, while at the highest dose used (20 mg/min) PGE2 induces renal vasoconstriction^[3].

PROTOCOL (Extracted from published papers and Only for reference)

Cell Assay: Prostaglandin E2 is dissolved in ethanol.^[1] Lymphocytes in CM (1×10⁶ cells/mL) are distributed in microculture plates (100 μL) in triplicate in the presence of PGE-treated T cells or medium-treated T cells and stimulated with PHA-P at various mitogenic doses. After 72 hr, cultures are pulsed with 1 μCi [³H]thymidine per well (specific activity 5 Ci/mM) for 16 to 18 hr, collected with a microprecipitator, dried, and counted in a liquid scintillation counter. **Animal Administration:** Prostaglandin E2 is dissolved in 95% ethanol and 0.002 M sodium carbonate and then diluted in phosphate buffered saline (PBS) to the desired concentration.^[2] Male Sprague Dawley rats (200-250 g) are used throughout the study. For 3 consecutive days rats in the experimental groups receive a daily intraperitoneal injection of either PGE2 (0.3 μg/kg body weight (BW)), the prostaglandin inhibitor meclofenamate (10 mg/kg BW) or the prostaglandin precursor arachidonic acid (0.3 μg/kg BW). To determine whether or not 0.3 μg/kg BW of a fatty acid produces nonspecific effects, the biologically inactive fatty acid 11, 14, 17-eicosatrienoic acid is also administered to a group of rats. Rats in the control group receive an equivalent volume (2.0 mL/kg BW) of the vehicle. On the third day, 3 mL of a suspension containing 1.2×10⁶ fluorescent methacrylate microbeads/mL of PBS are injected intraperitoneally (ip) into each rat. Six hours later all animals are given ip a regular dose of their respective treatment. Peritoneal exudate cells are harvested 19-22 hr later.

References:

[1]. Chouaib S, et al. The mechanisms of inhibition of human IL 2 production. II. PGE2 induction of suppressor T lymphocytes. J Immunol. 1984 Apr;132(4):1851-7.

[2]. Fernandez-Repollet E, et al. In vivo effects of prostaglandin E2 and arachidonic acid on phagocytosis of fluorescent methacrylate microbeads by rat peritoneal macrophages. J Histochem Cytochem. 1982 May;30(5):466-70.

[3]. Haylor J, et al. Renal vasodilator activity of prostaglandin E2 in the rat anaesthetized with pentobarbitone. Br J Pharmacol. 1982 May;76(1):131-7.

CAIndexNames:

Prosta-5,13-dien-1-oic acid, 11,15-dihydroxy-9-oxo-, (5Z,11 α ,13E,15S)-

SMILES:

CCCC[C@H](O)/C=C/[C@@H]1[C@H](C(C[C@H]1O)=O)C/C=C\CCCC(O)=O

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 732-484-9848 Fax: 888-484-5008 E-mail: sales@ChemScene.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA