

# **Product Data Sheet**

### Anti-PTP1B Antibody

Catalog #	Source	Reactivity	Applications
CPA4181	Rabbit	H, M, R, C, D, Mk, P, Z	WB, IH, IF/IC
Description	R	Rabbit polyclonal antibody to P	PTP1B
Immunogen	К	(LH-conjugated synthetic pepti	ide encompassing a sequence within the N-term
	re	egion of human PTP1B. The ex	act sequence is proprietary.
Purification	т	he antibody was purified by ir	nmunogen affinity chromatography.
Specificity	R	Recognizes endogenous levels o	of PTP1B protein.
Clonality	Р	Polyclonal	
Form	Li	iquid in 0.42% Potassium phos	sphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,
	а	and 0.01% sodium azide.	
Dilution	V	NB (1/500 - 1/1000), IH (1/100 -	1/200), IF/IC (1/100 - 1/500)
Gene Symbol	P	PTPN1	
Alternative N	ames P	PTP1B; Tyrosine-protein phosp	hatase non-receptor type 1; Protein-tyrosine
	р	bhosphatase 1B; PTP-1B	
Entrez Gene	5	5770 (Human); 19246 (Mouse);	; 24697 (Rat)
SwissProt	Р	218031 (Human); P35821 (Mou	use); P20417 (Rat)
Storage/Stabi	i <b>lity</b> S	hipped at 4°C. Upon delivery a	aliquot and store at -20°C for one year. Avoid
	fr	reeze/thaw cycles.	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**


WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com




For research purposes only, not for human use

# **Product Data Sheet**

Western blot analysis of PTP1B expression in HeLa (A), Jurkat (B) whole cell lysates.



Immunohistochemical analysis of PTP1B staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugad compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of PTP1B staining in SKNSH cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com