

Product Data Sheet

Anti-p53 (AcK319) Antibody

Catalog #	Source	Reactivity	Applications
CPA4292	Rabbit	H, B, Mk	WB, IH, IF/IC
Description	Rab	bit polyclonal antibody	to p53 (AcK319)
Immunogen	KLH	-conjugated synthetic p	eptide encompassing a sequence within the C-term
	regi	on of human p53 (AcK3	19). The exact sequence is proprietary.
Purification	The	antibody was purified I	by immunogen affinity chromatography.
Specificity	Rec	ognizes endogenous lev	els of p53 (AcK319) protein.
Clonality	Poly	yclonal	
Form	Liqu	սid in 0.42% Potassium լ	phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,
	and	0.01% sodium azide.	
Dilution	WB	(1/500 - 1/1000), IH (1/1	00 - 1/200), IF/IC (1/100 - 1/500)
Gene Symbol	TP5	3	
Alternative Na	ames P53	; Cellular tumor antiger	p53; Antigen NY-CO-13; Phosphoprotein p53; Tumor
	sup	pressor p53	
Entrez Gene	715	7 (Human)	
SwissProt	P04	637 (Human)	
Storage/Stabi	lity Ship	oped at 4°C. Upon delive	ery aliquot and store at -20°C for one year. Avoid
	free	eze/thaw cycles.	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com

Coherion

kDa


200

Δ


For research purposes only, not for human use

Product Data Sheet

Western blot analysis of p53 (AcK319) expression in HeLa TSA-treated (A) whole cell lysates.



Immunohistochemical analysis of p53 (AcK319) staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugad compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of p53 (AcK319) staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com