

Product Data Sheet

Anti-PI3K p85 alpha/p55 gamma (pY467/199) Antibody

Catalog #	Source	Reactivity	Applications		
CPA4491	Rabbit	H, M, R, B	WB, IH, IF/IC		
Description		Rabbit polyclonal antibody to PI3K p85 alpha/p55 gamma (pY467/199)			
Immunogen		KLH-conjugated synthetic peptide encompassing a sequence within the center			
		region of human PI3K p85 alpha/p55 gamma (pY467/199). The exact sequence is			
		proprietary.			
Purification	The antibody was purified by immunogen affinity chromatography.				
Specificity		Recognizes endogenous lev	vels of PI3K p85 alpha/p55 gamma (pY467/199) protein.		
Clonality		Polyclonal			
Form		Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,			
		and 0.01% sodium azide.			
Dilution		WB (1⁄500 - 1⁄1000), IH (1⁄1	00 - 1/200), IF/IC (1/100 - 1/500)		
Gene Symbol		PIK3R1; PIK3R3			
Alternative Names		PIK3R1; GRB1; Phosphatidy	linositol 3-kinase regulatory subunit alpha; PI3-kinase		
		regulatory subunit alpha; P	I3K regulatory subunit alpha; PtdIns-3-kinase regulatory		
		subunit alpha; Phosphatidy	linositol 3-kinase 85 kDa regulatory subunit alpha;		
		PI3-kinase subunit p85-alp	na; PtdIns-3-kinase regulatory subunit p85-alpha; PIK3R3;		
		Phosphatidylinositol 3-kina	se regulatory subunit gamma; PI3-kinase regulatory		
		subunit gamma; PI3K regul	atory subunit gamma; PtdIns-3-kinase regulatory subunit		
		gamma; Phosphatidylinosi	ol 3-kinase 55 kDa regulatory subunit gamma; PI3-kinase		
		subunit p55-gamma; PtdIn	s-3-kinase regulatory subunit p55-gamma; p55PIK		
Entrez Gene		5295 (Human); 18708, 187	10 (Mouse); 25513, 60664 (Rat)		
SwissProt		P27986, Q92569 (Human);	P26450, Q64143 (Mouse); Q63787, Q63789 (Rat)		
A sultanting large			histochowistwa IF, Jacob of Langerson of FC, Flow with working IC		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com




For research purposes only, not for human use

Product Data Sheet


Storage/Stability

Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid


freeze/thaw cycles.



Western blot analysis of PI3K p85 alpha/p55 gamma (pY467/199) expression in HEK293T (A), NIH3T3 (B) whole cell lysates.



Immunohistochemical analysis of PI3K p85 alpha/p55 gamma (pY467/199) staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugad compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of PI3K p85 alpha/p55 gamma (pY467/199) staining in HEK293T cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com