

## **Product Data Sheet**

### Anti-V5-tag Antibody

Source	Reactivity	Applications		
Mouse	N/A	WB, IF/IC, IP		
Мс	ouse monoclonal to V5	i-tag		
KLł	KLH-conjugated synthetic peptide encompassing a sequence of V5-tag. The exact			
sec	sequence is proprietary.			
Ree	Recognizes C-terminal, internal, and N-terminal V5-tag fusion proteins.			
Мс	Monoclonal			
Liq	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,			
and	d 0.01% sodium azide.			
WE	WB (1/2000 - 1/5000), IF/IC (1/200 - 1/500), IP (1/100 - 1/200)			
mes				
<b>ty</b> Shi	ipped at 4°C. Upon del	ivery aliquot and store at -20°C for one year. Avoid		
fre	eze/thaw cycles.			
	Mouse Mo KLI sed Mo Liq an WB	Mouse N/A Mouse monoclonal to V5 KLH-conjugated synthetic sequence is proprietary. Recognizes C-terminal, in Monoclonal Liquid in 0.42% Potassiur and 0.01% sodium azide. WB (1/2000 - 1/5000), IF/I		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**


WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



For research purposes only, not for human use

# **Product Data Sheet**

Western blot analysis of over-expressed V5-tagged protein in 293T cell lysate.



Immunofluorescent analysis of V5-tag staining in 293T cells transfected with a V5-tag protein. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Immunoprecipitation of V5-tagged protein from HEK293T cells transfected with vector overexpressing V5 tag, using Anti-V5-tag Antibody.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com