

# Product Data Sheet

## Anti-NPT2 Antibody

| Catalog #                | Source                                                                                                                                                                                                                                                          | Reactivity | Applications |
|--------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------|--------------|
| CQA3211                  | Rabbit                                                                                                                                                                                                                                                          | H, M, R    | WB, IF/IC    |
| <b>Description</b>       | Rabbit polyclonal antibody to NPT2                                                                                                                                                                                                                              |            |              |
| <b>Immunogen</b>         | Recombinant full length protein of human NPT2                                                                                                                                                                                                                   |            |              |
| <b>Purification</b>      | The antibody was purified by immunogen affinity chromatography.                                                                                                                                                                                                 |            |              |
| <b>Specificity</b>       | Recognizes endogenous levels of NPT2 protein.                                                                                                                                                                                                                   |            |              |
| <b>Clonality</b>         | Polyclonal                                                                                                                                                                                                                                                      |            |              |
| <b>Conjugation</b>       |                                                                                                                                                                                                                                                                 |            |              |
| <b>Form</b>              | Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.                                                                                                                                                       |            |              |
| <b>Dilution</b>          | WB (1/500 - 1/2000), IF/IC (1/50 - 1/200)                                                                                                                                                                                                                       |            |              |
| <b>Gene Symbol</b>       | SLC34A1                                                                                                                                                                                                                                                         |            |              |
| <b>Alternative Names</b> | NPT2; SLC17A2; Sodium-dependent phosphate transport protein 2A; Sodium-phosphate transport protein 2A; Na(+)-dependent phosphate cotransporter 2A; NaPi-3; Sodium/phosphate cotransporter 2A; Na(+)/Pi cotransporter 2A; NaPi-2a; Solute carrier family 34 memb |            |              |
| <b>Entrez Gene</b>       | 6569 (Human); 25548 (Rat)                                                                                                                                                                                                                                       |            |              |
| <b>SwissProt</b>         | Q06495 (Human); Q60825 (Mouse); Q06496 (Rat)                                                                                                                                                                                                                    |            |              |
| <b>Storage/Stability</b> | Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.                                                                                                                                                                |            |              |

**Application key:** E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

**Species reactivity key:** H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

### COHESION BIOSCIENCES LIMITED


**WEB**  
www.cohesionbio.com

**ORDER**  
order@cohesionbio.com


**SUPPORT**  
techsupport@cohesionbio.com

**CUSTOM**  
custom@cohesionbio.com

## Product Data Sheet



Western blot analysis of NPT2 expression in A549 (A), mouse kidney (B), rat lung (C) whole cell lysates.



Immunofluorescent analysis of NPT2 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

**Application key:** E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

**Species reactivity key:** H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

**COHESION BIOSCIENCES LIMITED**

**WEB**  
[www.cohesionbio.com](http://www.cohesionbio.com)

**ORDER**  
[order@cohesionbio.com](mailto:order@cohesionbio.com)

**SUPPORT**  
[techsupport@cohesionbio.com](mailto:techsupport@cohesionbio.com)

**CUSTOM**  
[custom@cohesionbio.com](mailto:custom@cohesionbio.com)